arXiv:1206.0439v1 [math-ph] 3 Jun 2012

From simplicial Chern-Simons theory to the shadow invariant I

March 6, 2013

Atle Hahn

Grupo de Física Matemática da Universidade de Lisboa
Av. Prof. Gama Pinto, 2
PT-1649-003 Lisboa, Portugal
Email: atle.hahn@gmx.de

Abstract

This is the first of a series of papers in which we develop a "discretization approach" for the rigorous realization of the non-Abelian Chern-Simons path integral for manifolds M of the form $M = \Sigma \times S^1$ and arbitrary simply-connected compact structure groups G. More precisely, we will introduce, for general links L in M, a rigorous version $\mathrm{WLO}_{rig}(L)$ of (the expectation values of) the corresponding Wilson loop observable $\mathrm{WLO}(L)$ in the so-called "torus gauge" by Blau and Thompson (Nucl. Phys. B408(1):345–390, 1993). For a simple class of links L we then evaluate $\mathrm{WLO}_{rig}(L)$ explicitly in a non-perturbative way, finding agreement with Turaev's shadow invariant $|L|$.

1 Introduction

In a celebrated paper, cf. [49], Witten succeeded in defining, on a physical level of rigor, a large class of new 3-manifold and link invariants ("Jones-Witten invariants") by making use of arguments based on the heuristic Chern-Simons path integral. Later, Reshetikhin and Turaev found a rigorous definition of these Jones-Witten invariants using the representation theory of quantum groups and suitable surgery operations on the base manifold, cf. [42, 41] and part I of [47]. A related approach is the so-called "shadow world" approach by Turaev, cf. [48] and part II of [47], which also works with quantum group representations but eliminates the use of surgery operations. In fact, the shadow invariant of a link is simply given by a finite "state sum" (see Appendix B in [28] for an "internal" reference in the special case relevant for us).

For several reasons (see, e.g., [36] and the introduction in [45]) it is desirable to find a rigorous "path integral definition" of the Jones-Witten invariants, i.e. a definition which is obtained by making rigorous sense of Witten's path integral expressions, either before or after a suitable gauge fixing has been applied.

Since this problem seems to be rather difficult it looks like a good strategy to restrict one's attention first to special base manifolds for which convenient gauge fixing procedures are available. A particularly promising situation is the case where the base manifold is of the form $M = \Sigma \times S^1$ and where we apply the so-called "torus gauge fixing" procedure, which was introduced by Blau and Thompson in [11] for the study of Chern-Simons models on such manifolds.

The results in [26, 27], which were obtained by extending and combining the work in [11, 12, 13, 24, 25] and [3, 23], suggest that for such manifolds it should be possible to find a rigorous realization of the heuristic path integral expressions for the Wilson loop observables (WLOs) which appear after torus gauge fixing has been applied, cf. Eq. (2.53) below.

Let us remark, however, that the approach in [26, 27] is quite technical[1]. It therefore makes sense to look for alternative and more elementary approaches. In the present paper and its sequel [28] we propose such an alternative approach, which was inspired by Adams' "simplicial" framework for Abelian Chern-Simons models, cf. [1, 2]. More precisely, we will develop two different but closely related approaches, "Approach I" and "Approach II" (cf. Remark 5.6 below and Sec. 3.4 in [28] for a short comparison of these two approaches). Our main result will be Theorem 5.3 in [28], which can be seen as a rigorous and more general version[2] of the main result in [11], cf. Sec. 7 in [11].

The present paper is organized as follows:

In Sec. 2 we summarize the relevant results of the torus gauge fixing approach of [11, 12, 13, 24, 25, 26] on Chern-Simons models on manifolds M of the form $M = \Sigma \times S^1$, following closely the presentation in [25]. At the end of this section we arrive at a heuristic formula for the WLOs, on which the rest of the present paper will be based, cf. Eq. (2.53).

In Sec. 3 we consider "oscillatory Gauss-type" (complex) measures on Euclidean vector spaces and discuss some of their properties.

In Sec. 4 we review the "simplicial" framework developed in [1, 2] for Abelian Chern-Simons theory.

In Sec. 5 we then present and discuss the first of the two aforementioned approaches for the discretization of the heuristic equation Eq. (2.53), i.e. "Approach I".

The present paper has an appendix consisting of six parts. In part A we list the Lie theoretic notation used in the present paper and we give some explicit formulas in the special case $G = SU(2)$. Part B is concerned with some technical issues arising in Sec. 2.2.4. In part C we give a formal definition of the notion of a "polyhedral cell decomposition" of a surface and the corresponding notion of "duality". Part D will be helpful for motivating the ansatz in Eq. (5.8) in Sec. 5.2 below. In part E we make some comments regarding the issue of continuum limits. Finally, in part F we give a brief summary of some aspects of the aforementioned rigorous continuum approach to CS theory on $M = \Sigma \times S^1$ in the torus gauge.

Note: In the present paper and in [28] we will work with general simply-connected compact Lie groups G since this does not involve much more work than we would have to invest if we restricted ourselves to a special case like, e.g., $G = SU(2)$. On the other hand treating the general situation makes it necessary to use several abstract concepts from Lie theory. The reader who prefers a more elementary treatment should feel free to concentrate on the special case $G = SU(2)$, cf. part A of the Appendix below for some useful formulas in this special case.

2 Chern-Simons theory on $M = \Sigma \times S^1$ in the torus gauge

2.1 Chern-Simons theory

Let us fix a simply-connected compact Lie group[3] G with Lie algebra $\mathfrak{g}$.

For every smooth manifolds X, every real vector space V and every $n \in \mathbb{N}$ we will denote by $\Omega^n(X, V)$ the space of V-valued n-forms on X and we set

$$\mathcal{A}_{X,V} := \Omega^1(X, V), \qquad \mathcal{A}_X := \mathcal{A}_{X,\mathfrak{g}} \tag{2.1}$$

By $\mathcal{G}_X$ we will denote the "gauge group" $C^\infty(X, G)$.

Let M be an oriented closed 3-manifold. In the following we will usually write $\mathcal{A}$ instead of $\mathcal{A}_M = \Omega^1(M, \mathfrak{g})$ and $\mathcal{G}$ instead of $\mathcal{G}_M$.

---

[1] see part F of the Appendix for a brief summary

[2] cf. Remark 5.4 in [28]

[3] cf. part A of the Appendix for concrete formulas in the special case $G = SU(2)$

For convenience let us consider for a while the special case where G is simple (cf. Remark 2.2 below for the case of general simply-connected compact Lie groups). The Chern-Simons action function $S_{CS} := S_{CS}(M, G, k)$ associated to M, G, and the "level" $k \in \mathbb{Z}\backslash\{0\}$ is then given by

$$S_{CS}(A) = -k\pi \int_M \langle A \wedge dA \rangle + \tfrac{1}{3} \langle A \wedge [A \wedge A] \rangle \in \mathbb{R}, \tag{2.2}$$

for all $A \in \mathcal{A}$. Above $[\cdot \wedge \cdot]$ denotes the wedge product associated to the bilinear map $[\cdot, \cdot] : \mathfrak{g} \times \mathfrak{g} \to \mathfrak{g}$ and where $\langle \cdot \wedge \cdot \rangle$ denotes the wedge product associated to the suitably[4] normalized Killing form $\langle \cdot, \cdot \rangle : \mathfrak{g} \times \mathfrak{g} \to \mathbb{R}$.

From the definition of S_{CS} it is obvious that S_{CS} is invariant under (orientation-preserving) diffeomorphisms. Thus, at a heuristic level, we can expect that the heuristic integral (the "partition function") $Z(M) := \int \exp(iS_{CS}(A))DA$ is a topological invariant of the 3-manifold M. Here DA denotes the informal "Lebesgue measure" on the space $\mathcal{A}$.

A similar statement holds if we consider (oriented and ordered) links L in M, i.e. finite tuples $L = (l_1, l_2, \ldots, l_m)$, $m \in \mathbb{N}$, where each l_i is a knot[5] in M such that $\operatorname{arc}(l_i) \cap \operatorname{arc}(l_j) = \emptyset$ holds whenever $i \neq j$. In the following we will identify each knot $l_i : S^1 \to M$ with the loop $[0, 1] \ni t \mapsto l_i(i_{S^1}(t)) \in M$ where $i_{S^1} : [0, 1] \ni s \mapsto \exp(2\pi i s) \in U(1) \cong S^1$.

If we fix a finite tuple $\rho = (\rho_1, \rho_2, \ldots, \rho_m)$ of finite-dimensional complex representations (="colors") of G then we can expect at a heuristic level that the mapping which maps every link $L = (l_1, l_2, \ldots, l_m)$ in M to the heuristic integral (the "expectation value of the Wilson loop observable associated to L and ρ")

$$\operatorname{WLO}(L, \rho) := \int_{\mathcal{A}} \prod_i \operatorname{Tr}_{\rho_i}\big(\operatorname{Hol}_{l_i}(A)\big) \exp(iS_{CS}(A))DA \tag{2.3}$$

is a link invariant. Here $\operatorname{Tr}_{\rho_i}$ denotes the trace in the representation ρ_i and $\operatorname{Hol}_{l_i}(A)$ denotes the holonomy of A around the loop l_i. Among the many different ways of writing $\operatorname{Hol}_{l_i}(A)$ explicitly the following equation will be particularly convenient for our purposes (cf. Sec. 5.3 below):

$$\operatorname{Hol}_{l_i}(A) = \lim_{n \to \infty} \prod_{k=1}^{n} \exp\left(\tfrac{1}{n} A\left(l_i'\left(\tfrac{k}{n}\right)\right)\right) \tag{2.4}$$

Here $\exp : \mathfrak{g} \to G$ is the exponential map of G.

Remark 2.1 We will simply write $\operatorname{WLO}(L)$ instead of $\operatorname{WLO}(L, \rho)$ if no confusion about the tuple $\rho = (\rho_1, \rho_2, \ldots, \rho_m)$ of "colors" can arise.

We remark that in the standard physics literature the notation $Z(M, L)$ is normally used instead of $\operatorname{WLO}(L)$.

For convenience we will assume (without loss of generality) in the following that the Lie group G fixed above is a Lie subgroup of $U(\mathbf{N})$, $\mathbf{N} \in \mathbb{N}$. The Lie algebra $\mathfrak{g}$ of G can then be identified with the obvious Lie subalgebra of the Lie algebra $u(\mathbf{N})$ of $U(\mathbf{N})$ and we have

$$\langle A, B \rangle = -\operatorname{Tr}(A \cdot B) \quad \forall A, B \in \mathfrak{g} \tag{2.5}$$

where "$\cdot$" denotes the matrix multiplication in $\operatorname{Mat}(\mathbf{N}, \mathbb{C})$ and where $\operatorname{Tr} := c \operatorname{Tr}_{\operatorname{Mat}(\mathbf{N}, \mathbb{C})}$ for suitably chosen[6] $c \in \mathbb{R}$. For example, in the special case $G = SU(\mathbf{N})$ we have $c = \frac{1}{4\pi^2}$.

---

[4] More precisely, the normalization is chosen such that $\langle \check{\alpha}, \check{\alpha} \rangle = 2$ for every short real coroot $\check{\alpha}$ w.r.t. any fixed Cartan subalgebra of $\mathfrak{g}$. Observe that after making the identification $\mathfrak{t} \cong \mathfrak{t}^*$ which is induced by $\langle \cdot, \cdot \rangle$ we have $\langle \alpha, \alpha \rangle = 2$ for every *long* root α. Thus the normalization here coincides with the one in [44]. This normalization guarantees that the exponential $\exp(iS_{CS})$ is "gauge invariant", i.e. invariant under the standard $\mathcal{G}$-operation on $\mathcal{A}$

[5] i.e. a smooth embedding $S^1 \to M$

[6] observe that if G is simple then every Ad-invariant scalar product on $\mathfrak{g}$ is proportional to the Killing form

The Chern-Simons action function S_{CS} can then be rewritten as

$$S_{CS}(A) = k\pi \int_M \mathrm{Tr}(A \wedge dA + \tfrac{2}{3} A \wedge A \wedge A), \quad A \in \mathcal{A} \tag{2.6}$$

where "$\wedge$" is now the wedge product for $(\mathrm{Mat}(\mathbf{N},\mathbb{C}), \cdot)$-valued forms. Moreover, on the RHS of Eq. (2.4) we can then reinterpret $\prod \cdots$ as the matrix product and exp as the exponential map of $\mathrm{Mat}(\mathbf{N},\mathbb{C})$.

Remark 2.2 Observe that a simply-connected compact Lie group is automatically semi-simple and can therefore be written as a product of the form $G = \prod_{i=1}^{r} G_i$, $r \in \mathbb{N}$, where each G_i is a simple simply-connected compact Lie group. We can generalize the definition of S_{CS} to this general situation by setting $S_{CS}(M, G, k)(A) := \sum_i S_{CS}(M, G_i, k)(A_i)$ for all $A \in \mathcal{A}$ where $(A_i)_i$ are the components of A w.r.t. to the decomposition $\mathfrak{g} = \oplus_i \mathfrak{g}_i$ ($\mathfrak{g}_i$ being the Lie algebra of G_i).

In view of Sec. 7 in [28] let us generalize the definition of S_{CS} even further and introduce for every sequence $(k_i)_{i \leq r}$ of non-zero integers the function $S_{CS}(M, G, (k_i)_i)$by setting $S_{CS}(M, G, (k_i)_i)(A) := \sum_i S_{CS}(M, G_i, k_i)(A_i)$ for all $A \in \mathcal{A}$. In fact, in the present paper and in [28] only two special cases will play a role, namely the case $r = 1$ (i.e. G simple) and the case $r = 2$, $G_2 = G_1$ and $k_2 = -k_1$, cf. Sec. 7 in [28].

2.2 Torus gauge fixing

For the rest of this paper let us fix a maximal torus T of G. The Lie algebra of T will be denoted by $\mathfrak{t}$.

2.2.1 Motivation

In order to motivate the definition of the torus gauge fixing procedure for the manifold M of the form $M = \Sigma \times S^1$ where Σ is a connected surface let us first have a quick look at the orbit space $\mathcal{A}/\mathcal{G}$ for the three manifolds $M = \mathbb{R}$, $M = S^1$, and $M = \Sigma \times \mathbb{R}$.

In the following $\frac{\partial}{\partial t}$ will denote the vector field on $\mathbb{R}$ (resp. S^1) which is induced by the map $\mathrm{id}_{\mathbb{R}}$ (resp. the map $i_{S^1} : [0,1] \ni s \mapsto \exp(2\pi i s) \in U(1) \cong S^1$) and dt will denote the dual 1-form on $\mathbb{R}$ (resp. S^1). The obvious "lift"/pullback of $\frac{\partial}{\partial t}$ and dt to the product manifolds $\Sigma \times \mathbb{R}$ and $\Sigma \times S^1$ will again be denoted by $\frac{\partial}{\partial t}$ and dt.

i) $M = \mathbb{R}$: Here every 1-form $A = A_0 dt \in \mathcal{A}$ is gauge-equivalent to the trivial 1-form $0\, dt = 0$, so $\mathcal{A}/\mathcal{G}$ has just one element.

ii) $M = S^1$: Here every $A \in \mathcal{A}$ is gauge equivalent[7] to a 1-form of the form Bdt with constant $B : S^1 \to \mathfrak{g}$. Moreover, according to the fundamental theorem of maximal tori we can choose B to be $\mathfrak{t}$-valued, so the map $\pi_{\mathcal{G}} : \{Bdt \mid B \in C^\infty(S^1, \mathfrak{t}) \text{ is constant } \} \to \mathcal{A}/\mathcal{G}$ is surjective.

iii) $M = \Sigma \times \mathbb{R}$: Every $A \in \mathcal{A}$ can be written uniquely in the form $A = A^\perp + A_0 dt$ with $A^\perp \in \mathcal{A}^\perp := \{A \in \mathcal{A} \mid A(\frac{\partial}{\partial t}) = 0\}$ and $A_0 \in C^\infty(M, \mathfrak{g})$. Using the argument for the case $M = \mathbb{R}$ for each of the "fibers" $\{\sigma\} \times \mathbb{R} \cong \mathbb{R}$, $\sigma \in \Sigma$, we can easily conclude that every 1-form A can be gauge-transformed into an element of $\mathcal{A}^\perp$. In other words, the map $\pi_{\mathcal{G}} : \mathcal{A}^\perp \to \mathcal{A}/\mathcal{G}$ is surjective.

After these preparations let us now go back to the original manifold.

---

[7] this follows, e.g., by looking at the explicit form of the well-known bijection $\psi : \mathcal{A}_{flat}/\mathcal{G} \to \mathrm{Hom}(\pi_1(M), G)/G$ and taking into account that in the special case $M = S^1$ we have $\mathcal{A}/\mathcal{G} = \mathcal{A}_{flat}/\mathcal{G}$ and $\mathrm{Hom}(\pi_1(M), G)/G \cong G/G$

iv) $M = \Sigma \times S^1$: Again every $A \in \mathcal{A}$ can be written uniquely in the form $A = A^\perp + A_0 dt$ with $A^\perp \in \mathcal{A}^\perp$ and $A_0 \in C^\infty(M, \mathfrak{g})$ where $\mathcal{A}^\perp$ is defined again by

$$\mathcal{A}^\perp := \{A \in \mathcal{A} \mid A(\tfrac{\partial}{\partial t}) = 0\} \tag{2.7}$$

Combining the results for the case $M = S^1$ and $M = \Sigma \times \mathbb{R}$ and making the identification $\{B \in C^\infty(\Sigma \times S^1, \mathfrak{t}) \mid B(\sigma, \cdot) \text{ constant for all } \sigma \in \Sigma\} \cong C^\infty(\Sigma, \mathfrak{t})$ one is naturally led to the space

$$\mathcal{A}^{qax}(T) := \mathcal{A}^\perp \oplus \{B dt \mid B \in C^\infty(\Sigma, \mathfrak{t})\} \tag{2.8}$$

and to the question whether the map $\pi_{\mathcal{G}} : \mathcal{A}^{qax}(T) \to \mathcal{A}/\mathcal{G}$ is surjective. For technical reasons let us also introduce the space

$$\mathcal{A}^{qax} := \mathcal{A}^\perp \oplus \{B dt \mid B \in C^\infty(\Sigma, \mathfrak{g})\} \tag{2.9}$$

In Sec. 2.2.4 we will study the map $\pi_{\mathcal{G}} : \mathcal{A}^{qax}(T) \to \mathcal{A}/\mathcal{G}$ in the situation relevant for us. Before we do this we will have to make a short digression where we introduce the two heuristic concepts of a generalized gauge fixing and an abstract gauge fixing which will be useful in Sec. 2.2.3 and Sec. 2.2.4 below.

2.2.2 Two heuristic concepts: "generalized" and "abstract" gauge fixing

Let us call a (not necessarily linear) subspace V of $\mathcal{A}$ a *gauge fixing subspace* iff its elements form a complete and minimal set of representatives of $\mathcal{A}/\mathcal{G}$, or, equivalently, iff the map $\Pi_V : V \to \mathcal{A}/\mathcal{G}$ which is obtained by restricting the canonical projection $\pi_{\mathcal{G}} : \mathcal{A} \to \mathcal{A}/\mathcal{G}$ onto V is a bijection. (Thus $\Pi_V^{-1} : \mathcal{A}/\mathcal{G} \to \mathcal{A}$ will be a *gauge fixing* in the usual sense).

Let $d\nu_{\mathcal{A}/\mathcal{G}}$ denote the image of the informal Lebesgue measure DA under $\pi_{\mathcal{G}}$, i.e.

$$d\nu_{\mathcal{A}/\mathcal{G}} := (\pi_{\mathcal{G}})_* DA \tag{2.10}$$

If V is a gauge fixing subspace then, setting $d\mu_V := (\Pi_V)_*^{-1}(d\nu_{\mathcal{A}/\mathcal{G}})$, we trivially have (at an informal level)

$$(\Pi_V)_* d\mu_V = d\nu_{\mathcal{A}/\mathcal{G}} \tag{2.11}$$

and therefore[8] also (informally)

$$\int \chi(A) DA = \int_V \chi(A) d\mu_V(A) \tag{2.12}$$

for every $\mathcal{G}$-invariant function $\chi : \mathcal{A} \to \mathbb{C}$.

Remark 2.3 If V is a "sufficiently nice" subspace of $\mathcal{A}$ the informal measure $d\mu_V$ will have an explicitly computable "density" w.r.t. to $DA_{|V}$. This density is usually called the "Faddeev-Popov determinant" (of the gauge fixing associated to V), cf., e.g., [40] and Appendix C in [26].

We will call a subset $V \subset \mathcal{A}$ a *generalized gauge fixing subspace* iff the map $\Pi_V : V \to \mathcal{A}/\mathcal{G}$ given as above is "essentially surjective" in the sense that the complement of $\pi_{\mathcal{G}}(V)$ in $\mathcal{A}/\mathcal{G}$ is a $d\nu_{\mathcal{A}/\mathcal{G}}$-zero subset. Since for such a generalized gauge fixing space the map $\Pi_V : V \to \mathcal{A}/\mathcal{G}$ need not be injective we can not hope to obtain an informal measure $d\mu_V$ fulfilling Eq. (2.11) above in a canonical way as above. However, since by assumption Π_V is essentially surjective there will be (non-canonical informal) measures $d\mu_V$ on V such that Eq. (2.11) above is fulfilled at an informal level. We will call any such[9] pair $(V, d\mu_V)$ a *generalized gauge fixing*.

---

[8] since $\int_{\mathcal{A}} \chi(A) DA = \int_{\mathcal{A}/\mathcal{G}} \bar{\chi}\, d\nu_{\mathcal{A}/\mathcal{G}} = \int_{\mathcal{A}/\mathcal{G}} \bar{\chi}\, (\Pi_V)_* d\mu_V = \int_V (\bar{\chi} \circ \Pi_V) d\mu_V = \int_V \chi(A) d\mu_V(A)$ where $\bar{\chi} : \mathcal{A}/\mathcal{G} \to \mathbb{C}$ is uniquely given by $\chi = \bar{\chi} \circ \pi_{\mathcal{G}}$

An *abstract gauge fixing* is a triple $(V, \Pi_V, d\mu_V)$ where V is an arbitrary (heuristic) measurable space, Π_V is a (heuristic) measurable map $V \to \mathcal{A}/\mathcal{G}$ and $d\mu_V$ a measure on V such that Eq. (2.11) above (and therefore also Eq. (2.12)) is fulfilled[10]. Of course, for an abstract gauge fixing to be useful the space V should have enough structure so that one can perform concrete computations[11]. For the abstract gauge fixing which will appear in Sec. 2.2.4 the space V will be the direct product of a linear space and a countable discrete space.

2.2.3 Torus gauge fixing for non-compact $M = \Sigma \times S^1$

In the following let $\mathfrak{k}$ denote the $\langle \cdot, \cdot \rangle$-orthogonal complement of $\mathfrak{t}$ in $\mathfrak{g}$ and set

$$G_{reg} := \{ g \in G \mid g \text{ is regular} \} \tag{2.13}$$

Recall that an element g of G is called "regular" iff it is contained in exactly one maximal torus of G, cf. part A of the Appendix below for concrete formulas in the special case $G = SU(2)$.

Let us now restrict to the special situation which is relevant for CS theory, i.e. where G is simply-connected, cf. Sec. 2.1. It turns out that if Σ is non-compact we have (cf. point ii) in part B of the Appendix)

$$\mathcal{A}_{reg}/\mathcal{G} \subset \mathcal{A}^{qax}(T)/\mathcal{G} = \pi_{\mathcal{G}}(\mathcal{A}^{qax}(T)) \tag{2.14}$$

where

$$\mathcal{A}_{reg} := \{ A \in \mathcal{A} \mid \mathrm{Hol}_{l_\sigma}(A) \in G_{reg} \text{ for every } \sigma \in \Sigma \} \tag{2.15}$$

and where l_σ is the "vertical" loop "above" σ, i.e. $l_\sigma : [0,1] \ni s \mapsto (\sigma, i_{S^1}(s)) \in M$ with i_{S^1} as above. Moreover, since $\mathrm{codim}(G \backslash G_{reg}) \geq 3$ (cf., e.g., Chap. V in [15]) and $\dim(\Sigma) = 2$, a "generic" function $f : \Sigma \to G$ will remain inside G_{reg}. Thus we can argue at a heuristic level that the difference sets

$$C^\infty(\Sigma, G) \backslash C^\infty(\Sigma, G_{reg}), \quad \mathcal{A} \backslash \mathcal{A}_{reg}, \quad (\mathcal{A}/\mathcal{G}) \backslash (\mathcal{A}_{reg}/\mathcal{G})$$

are all "negligible", i.e. "zero-subsets" w.r.t. $D\Omega$ (the heuristic Haar measure on $C^\infty(\Sigma, G)$) resp. DA resp. $d\nu_{\mathcal{A}/\mathcal{G}}$. Thus, if Σ is non-compact, the space $V := \mathcal{A}^{qax}(T)$ should indeed be a generalized gauge fixing space and there should be a measures $d\mu_V$ on V fulfilling Eq. (2.11).

In order to describe such a measure $d\mu_V$ explicitly, let us make the identification

$$V = \mathcal{A}^{qax}(T) \cong A^\perp \times \mathcal{B} \tag{2.16}$$

where we have set

$$\mathcal{B} := C^\infty(\Sigma, \mathfrak{t}) \tag{2.17}$$

Below $DA^\perp$ will denote the (informal) "Lebesgue measure" on $\mathcal{A}^\perp$ and DB the (informal) "Lebesgue measure" on $\mathcal{B}$.

It can be shown at an informal level[12] that the measure $d\mu_V$ on $V \cong A^\perp \times \mathcal{B}$ given by

$$d\mu_V := DA^\perp \otimes \left(\det\left(1_{\mathfrak{k}} - \exp(\mathrm{ad}(B))_{|\mathfrak{k}}\right) DB \right) \tag{2.18}$$

---

[9] i.e. V is a generalized gauge fixing space and $d\mu_V$ a measure on V fulfilling Eq. (2.11)

[10] in this case, one can conclude at an informal level that Π_V should then be essentially surjective in the sense above since one can argue that the informal measure $d\nu_{\mathcal{A}/\mathcal{G}}$ on $\mathcal{A}/\mathcal{G}$ should have "full support"

[11] this excludes, for example, the "trivial" abstract gauge fixing $V = \mathcal{A}/\mathcal{G}$, $\Pi_V = \mathrm{id}_{\mathcal{A}/\mathcal{G}}$ and $d\mu_V = d\nu_{\mathcal{A}/\mathcal{G}}$. The orbit space $\mathcal{A}/\mathcal{G}$ has so little structure that it is not very useful for explicit computations, which is exactly the reason why one usually tries to apply a suitable gauge fixing (in the usual or generalized sense)

[12] by computing the Faddeev-Popov determinant of a closely related (proper) gauge fixing, cf. Sec. 2.3 and Sec. 2.4 in [26] (and Remark 2.3 above). Observe that Eq. (2.19) below is just the analogue of Eq. (2.23) in Sec. 2.4 in [26] where P is replaced by $\mathfrak{t}$

fulfills Eq. (2.11) up to a multiplicative constant, or, equivalently, that

$$\int_{\mathcal{A}} \chi(A) DA \sim \int_{\mathcal{B}} \Big[\int_{\mathcal{A}^\perp} \chi(A^\perp + B dt) DA^\perp \Big] \det\big(1_{\mathfrak{k}} - \exp(\mathrm{ad}(B))_{|\mathfrak{k}}\big) DB \tag{2.19}$$

holds for every $\mathcal{G}$-invariant function $\chi : \mathcal{A} \to \mathbb{C}$.

Convention 1 *Above and in the following "$\sim$" denotes equality up to a multiplicative constant. Of course, this "constant" can/should depend on G and M but it is clearly independent of χ.*

2.2.4 Torus gauge fixing for compact $M = \Sigma \times S^1$

The case of compact Σ, which is the case we are actually interested in, requires more care since in this case we have $\mathcal{A}_{reg}/\mathcal{G} \not\subset \pi_{\mathcal{G}}(\mathcal{A}^{qax}(T))$ (cf. point iii) in part B of the Appendix; cf. also Example 2 in Sec. 3 in [13]). If one still wants to transform a general 1-form $A \in \mathcal{A}_{reg}$ into an element $A^\perp + B dt$ of $\mathcal{A}^{qax}(T)$ one can do so only if one uses a certain (mildly) singular gauge transformation Ω and also allows $A^\perp$ to have a similar singularity (cf. the maps Ω_{h} and the 1-forms $A_{\mathrm{sg}}(\mathrm{h})$ appearing in Step 2 and Step 3 below). For functions χ fulfilling an additional property (cf. Step 2 below) this strategy indeed allows us to generalize Eq. (2.19) so that also the case of compact surfaces Σ is included, cf. Eq (2.28) below.

We will only sketch the derivation of Eq (2.28). For more details we refer to [25] where a detailed derivation of a very similar equation is given where $\mathcal{B} = C^\infty(\Sigma, \mathfrak{t})$ is replaced by $C^\infty(\Sigma, P)$, $P \subset \mathfrak{t}$ being a fixed Weyl alcove.

Preparation for Step 1: Let us set[13]

$$[\Sigma, G/T] := C^\infty(\Sigma, G/T)/\mathcal{G}_\Sigma \tag{2.20}$$

where the expression on the RHS denotes the orbit space of the $\mathcal{G}_\Sigma$-operation on $C^\infty(\Sigma, G/T)$ given by $\bar{g} \cdot \Omega := \Omega^{-1}\bar{g}$ for $\bar{g} \in C^\infty(\Sigma, G/T)$ and $\Omega \in \mathcal{G}_\Sigma$. For each $\mathrm{h} \in [\Sigma, G/T]$ we pick a representative $\bar{g}_{\mathrm{h}} \in C^\infty(\Sigma, G/T)$. We will keep each $\bar{g}_{\mathrm{h}}$ fixed in the following.

For each $\bar{g} \in G/T$ and $b \in \mathfrak{t}$ we set $\bar{g} b \bar{g}^{-1} := g b g^{-1} \in \mathfrak{g}$ where g is an arbitrary[14] element of G fulfilling $gT = \bar{g}$.

Step 1: We now introduce a suitable abstract gauge fixing $(\overline{V}, \Pi_{\overline{V}}, d\mu_{\overline{V}})$ ("abstract torus gauge fixing"). We take

$$\overline{V} := \mathcal{A}^\perp \times \mathcal{B} \times [\Sigma, G/T] \tag{2.21}$$

and define $\Pi_{\overline{V}} : \overline{V} \to \mathcal{A}/\mathcal{G}$ by

$$\Pi_{\overline{V}}(A^\perp, B, \mathrm{h}) = \pi_{\mathcal{G}}(A^\perp + \bar{g}_{\mathrm{h}} B \bar{g}_{\mathrm{h}}^{-1} dt) \tag{2.22}$$

for all $(A^\perp, B, \mathrm{h}) \in \overline{V}$ where $\bar{g}_{\mathrm{h}} B \bar{g}_{\mathrm{h}}^{-1} \in C^\infty(\Sigma, \mathfrak{g})$ is given by $(\bar{g}_{\mathrm{h}} B \bar{g}_{\mathrm{h}}^{-1})(\sigma) := \bar{g}_{\mathrm{h}}(\sigma) B(\sigma) \bar{g}_{\mathrm{h}}^{-1}(\sigma)$ for all $\sigma \in \Sigma$. According to point i) in part B of the Appendix we have

$$\mathcal{A}_{reg}/\mathcal{G} \subset \mathrm{Image}(\Pi_{\overline{V}}), \tag{2.23}$$

so $\Pi_{\overline{V}}$ is essentially surjective in the sense above. Thus we can hope to be able to find a heuristic measure $d\mu_{\overline{V}}$ on $\overline{V}$ fulfilling Eq. (2.11) (with V replaced by $\bar{V}$). Since $[\Sigma, G/T]$ is a countable set (cf. Remark 2.5 below) it is tempting to try the ansatz

$$d\mu_{\overline{V}} := DA^\perp \otimes \big(\det\big(1_{\mathfrak{k}} - \exp(\mathrm{ad}(B))_{|\mathfrak{k}}\big) DB\big) \otimes \# \tag{2.24}$$

---

[13] the notation $[\Sigma, G/T]$ is motivated by the fact that $C^\infty(\Sigma, G/T)/\mathcal{G}_\Sigma$ coincides with the set of homotopy classes of maps $\Sigma \to G/T$, cf. Proposition 3.2 in [25]

[14] observe that for $b \in \mathfrak{t}$ the value $\bar{g} b \bar{g}^{-1}$ will not depend on the special choice of g

where # is the counting measure on $[\Sigma, G/T]$.

It turns out that with this choice Eq. (2.11) is indeed fulfilled at a heuristic level (with V replaced by $\bar{V}$ and with "=" replaced by "$\sim$"), or, equivalently, that we have

$$\int_{\mathcal{A}} \chi(A) DA$$
$$\sim \sum\nolimits_{\mathrm{h} \in [\Sigma, G/T]} \int_{\mathcal{B}} \Big[\int_{\mathcal{A}^{\perp}} \chi(A^{\perp} + \bar{g}_{\mathrm{h}} B \bar{g}_{\mathrm{h}}^{-1} dt) DA^{\perp} \Big] \det\bigl(1_{\mathfrak{k}} - \exp(\mathrm{ad}(B))_{|\mathfrak{k}}\bigr) DB \quad (2.25)$$

for every $\mathcal{G}$-invariant function $\chi : \mathcal{A} \to \mathbb{C}$, cf. Eq. (2.26) in [26].

Observe that the function $\bar{g}_{\mathrm{h}} B \bar{g}_{\mathrm{h}}^{-1} \in C^{\infty}(\Sigma, \mathfrak{g})$ will in general not take only values in $\mathfrak{t}$ unless $\mathrm{h} = [1_T]$ (here 1_T is the constant function on Σ which only takes the value $T \in G/T$). This reduces the usefulness of Eq. (2.25) considerably. Fortunately, for a certain class of functions χ one can derive a more useful variant of Eq. (2.25), cf. Eq. (2.28) below.

Preparation for Step 2: Let us fix a point $\sigma_0 \in \Sigma$ and pick, for each $\mathrm{h} \in [\Sigma, G/T]$, a lift $\Omega_{\mathrm{h}} \in \mathcal{G}_{\Sigma \backslash \{\sigma_0\}} = C^{\infty}(\Sigma \backslash \{\sigma_0\}, G)$ of $(\bar{g}_{\mathrm{h}})_{|\Sigma \backslash \{\sigma_0\}} \in C^{\infty}(\Sigma \backslash \{\sigma_0\}, G/T)$, cf. Remark 2.4 below. We will keep σ_0 and each Ω_{h} fixed for the rest of this paper.

Remark 2.4 The existence of the "lifts" $\Omega_{\mathrm{h}} \in \mathcal{G}_{\Sigma \backslash \{\sigma_0\}}$ of $(\bar{g}_{\mathrm{h}})_{|\Sigma \backslash \{\sigma_0\}}$ picked above is guaranteed by the following three observations:

i) the surface $\Sigma \backslash \{\sigma_0\}$ (and in fact every non-compact surface) is homotopy equivalent to a 1-complex

ii) G/T is simply-connected

iii) $\pi_{G/T} : G \to G/T$ is a fiber bundle and therefore possesses the homotopy lifting property (cf., e.g., [32])

The first two observations imply that each of the maps $(\bar{g}_{\mathrm{h}})_{|\Sigma \backslash \{\sigma_0\}}$ is 0-homotopic and the third observation then implies the existence of Ω_{h} (cf. also Example B.1 in part B of the Appendix below).

Clearly, the restriction mapping $\mathcal{G}_{\Sigma} \ni \Omega \mapsto \Omega_{|\Sigma \backslash \{\sigma_0\}} \in \mathcal{G}_{\Sigma \backslash \{\sigma_0\}}$ is injective so we can consider $\mathcal{G}_{\Sigma}$ to be a subgroup of $\mathcal{G}_{\Sigma \backslash \{\sigma_0\}}$. In a similar way, we will identify $C^{\infty}(\Sigma, \mathfrak{g})$ with a subspace of $C^{\infty}(\Sigma \backslash \{\sigma_0\}, \mathfrak{g})$, $\mathcal{A}^{\perp}$ with a subspace of $\mathcal{A}^{\perp}_{(\Sigma \backslash \{\sigma_0\}) \times S^1}$ (defined in the obvious way), and $\mathcal{A}^{qax} = \mathcal{A}^{\perp} \oplus C^{\infty}(\Sigma, \mathfrak{g}) dt$ with a subspace of $\mathcal{A}^{qax}_{(\Sigma \backslash \{\sigma_0\}) \times S^1} := \mathcal{A}^{\perp}_{(\Sigma \backslash \{\sigma_0\}) \times S^1} \oplus C^{\infty}(\Sigma \backslash \{\sigma_0\}, \mathfrak{g}) dt$. Then we have

$$\begin{array}{rcl}
\mathcal{G}_{\Sigma} \subset & \overline{\mathcal{G}_{\Sigma}} := \langle \mathcal{G}_{\Sigma} \cup \{\Omega_{\mathrm{h}} \mid \mathrm{h} \in [\Sigma, G/T]\} \rangle & \subset \mathcal{G}_{\Sigma \backslash \{\sigma_0\}} \\
C^{\infty}(\Sigma, \mathfrak{g}) \subset & \overline{C^{\infty}}(\Sigma, \mathfrak{g}) := \{B \in C^{\infty}(\Sigma \backslash \{\sigma_0\}, \mathfrak{g}) \mid B \text{ bounded } \} & \subset C^{\infty}(\Sigma \backslash \{\sigma_0\}, \mathfrak{g}) \\
\mathcal{A}^{\perp} \subset & \overline{\mathcal{A}^{\perp}} := \mathcal{A}^{\perp} \cdot \overline{\mathcal{G}_{\Sigma}} & \subset \mathcal{A}^{\perp}_{(\Sigma \backslash \{\sigma_0\}) \times S^1} \\
\mathcal{A}^{qax} \subset & \overline{\mathcal{A}^{qax}} := \overline{\mathcal{A}^{\perp}} \oplus \overline{C^{\infty}}(\Sigma, \mathfrak{g}) dt & \subset \mathcal{A}^{qax}_{(\Sigma \backslash \{\sigma_0\}) \times S^1}
\end{array}$$

Observe that the spaces $\overline{C^{\infty}}(\Sigma, \mathfrak{g})$, $\overline{\mathcal{A}^{\perp}}$, and $\overline{\mathcal{A}^{qax}}$ are $\overline{\mathcal{G}_{\Sigma}}$-invariant subsets of the spaces $C^{\infty}(\Sigma \backslash \{\sigma_0\}, \mathfrak{g})$ resp. $\mathcal{A}^{\perp}_{(\Sigma \backslash \{\sigma_0\}) \times S^1}$ resp. $\mathcal{A}^{qax}_{(\Sigma \backslash \{\sigma_0\}) \times S^1}$ (when the latter are equipped with the obvious $\mathcal{G}_{\Sigma \backslash \{\sigma_0\}}$-operation).

Step 2: Let $\chi : \mathcal{A} \to \mathbb{C}$ be a $\mathcal{G}$-invariant function with the extra property that the function $\chi^{qax} : \mathcal{A}^{qax} \to \mathbb{C}$ given by $\chi^{qax} := \chi_{|\mathcal{A}^{qax}}$ is not only $\mathcal{G}_{\Sigma}$-invariant but can also be extended to a $\overline{\mathcal{G}_\Sigma}$-invariant function $\overline{\chi^{qax}} : \overline{\mathcal{A}^{qax}} \to \mathbb{C}$. If χ has this property we obtain for the integrand in the inner integral on the right-hand side of (2.25)

$$\begin{aligned}\chi(A^\perp + (\bar{g}_\mathrm{h} B \bar{g}_\mathrm{h}^{-1})dt) = \chi^{qax}(A^\perp + (\bar{g}_\mathrm{h} B \bar{g}_\mathrm{h}^{-1})dt) = \overline{\chi^{qax}}(A^\perp + (\Omega_\mathrm{h} B \Omega_\mathrm{h}^{-1})dt)\\ = \overline{\chi^{qax}}((A^\perp \cdot \Omega_\mathrm{h}) + Bdt) = \overline{\chi^{qax}}(\Omega_\mathrm{h}^{-1} A^\perp \Omega_\mathrm{h} + \Omega_\mathrm{h}^{-1} d\Omega_\mathrm{h} + Bdt) \quad (2.26)\end{aligned}$$

So for such a function χ we arrive at the following modification of Eq. (2.25)

$$\int_\mathcal{A} \chi(A) DA \sim \sum\nolimits_{\mathrm{h} \in [\Sigma, G/T]} \int_\mathcal{B} \Big[\int_{\mathcal{A}^\perp} \overline{\chi^{qax}}(\Omega_\mathrm{h}^{-1} A^\perp \Omega_\mathrm{h} + \Omega_\mathrm{h}^{-1} d\Omega_\mathrm{h} + Bdt) DA^\perp \Big] \\ \times \det\big(1_\mathfrak{k} - \exp(\mathrm{ad}(B))_{|\mathfrak{k}}\big) DB \quad (2.27)$$

Step 3: As a final step let us simplify Eq. (2.27) by the change of variable $\Omega_\mathrm{h}^{-1} A^\perp \Omega_\mathrm{h} + \pi_\mathfrak{k}(\Omega_\mathrm{h}^{-1} d\Omega_\mathrm{h}) \to A^\perp$ where $\pi_\mathfrak{k}$ is the $\langle \cdot, \cdot \rangle$-orthogonal projection $\mathfrak{g} \to \mathfrak{k}$.

For the special χ relevant for us (cf. Sec. 2.3.1 below), this change of variable can indeed be justified[15], cf. Sec. 4.2 in [25]. Applying this change of variable to Eq. (2.27) we finally obtain

$$\int_\mathcal{A} \chi(A) DA \sim \sum\nolimits_{\mathrm{h} \in [\Sigma, G/T]} \int_\mathcal{B} \Big[\int_{\mathcal{A}^\perp} \overline{\chi^{qax}}(A^\perp + A_\mathrm{sg}(\mathrm{h}) + Bdt) DA^\perp \Big] \\ \times \det\big(1_\mathfrak{k} - \exp(\mathrm{ad}(B))_{|\mathfrak{k}}\big) DB \quad (2.28)$$

where we have set

$$A_\mathrm{sg}(\mathrm{h}) := \pi_\mathfrak{t}(\Omega_\mathrm{h}^{-1} d\Omega_\mathrm{h}) \quad (2.29)$$

where $\pi_\mathfrak{t}$ is the $\langle \cdot, \cdot \rangle$-orthogonal projection $\mathfrak{g} \to \mathfrak{t}$.

Observe that we can equally well work with the abstract gauge fixing space $\overline{V} := \mathcal{A}^\perp \times C^\infty(\Sigma, \mathfrak{t}_{reg}) \times [\Sigma, G/T]$ instead of $\overline{V} := \mathcal{A}^\perp \times \mathcal{B} \times [\Sigma, G/T]$ in Step 1 above where

$$\mathfrak{t}_{reg} := \exp^{-1}(T_{reg}) \quad (2.30)$$

with $T_{reg} := T \cap G_{reg}$ ($\mathfrak{t}_{reg}$ is just the union of all the Weyl alcoves).

If we do so we arrive at[16]

$$\int_\mathcal{A} \chi(A) DA \sim \sum\nolimits_{\mathrm{h} \in [\Sigma, G/T]} \int_{C^\infty(\Sigma, \mathfrak{t}_{reg})} \Big[\int_{\mathcal{A}^\perp} \overline{\chi^{qax}}(A^\perp + A_\mathrm{sg}(\mathrm{h}) + Bdt) DA^\perp \Big] \\ \times \det\big(1_\mathfrak{k} - \exp(\mathrm{ad}(B))_{|\mathfrak{k}}\big) DB \quad (2.31)$$

which will be slightly more convenient in Sec. 2.3.1 below.

Remark 2.5 The mapping $n : [\Sigma, G/T] \to \mathfrak{t}$ given by $n(\mathrm{h}) = \int_{\Sigma \backslash \{\sigma_0\}} dA_\mathrm{sg}(\mathrm{h})$ is independent of the special choice of $\bar{g}_\mathrm{h}$ and Ω_h involved. Moreover, one can show that n is a bijection from $[\Sigma, G/T]$ onto the lattice

$$I := \ker(\exp_{|\mathfrak{t}}) \quad (2.32)$$

(see Proposition 3.5, Proposition 3.6, and Remark 3.2 in [25]; cf. also Sec. 5 in [13] and Example B.1 in part B of the Appendix below)

---

[15] by contrast the more radical change of variable $\Omega_\mathrm{h}^{-1} A^\perp \Omega_\mathrm{h} + \Omega_\mathrm{h}^{-1} d\Omega_\mathrm{h} \to A^\perp$ is not admissible (at least not in the present form), cf. Remark 2.6 below

[16] alternatively, one could try to "derive" Eq. (2.31) directly form Eq. (2.28) by argueing that, since $\det\big(1_\mathfrak{k} - \exp(\mathrm{ad}(b))_{|\mathfrak{k}}\big) = 0$ for every $b \in \mathfrak{t} \backslash \mathfrak{t}_{reg}$ the set $C^\infty(\Sigma, \mathfrak{t}) \backslash C^\infty(\Sigma, \mathfrak{t}_{reg})$ should be a $\det\big(1_\mathfrak{k} - \exp(\mathrm{ad}(B))_{|\mathfrak{k}}\big) DB$-zero subset of $\mathcal{B} = C^\infty(\Sigma, \mathfrak{t})$

2.3 Chern-Simons theory in the torus gauge

2.3.1 Application of Eq. (2.31) to Chern-Simons theory

From now on we will assume that the (connected) surface Σ fixed above is oriented and compact. Let $L = ((l_1, l_2, \ldots, l_m), (\rho_1, \rho_2, \ldots, \rho_m))$ be a colored link in $M = \Sigma \times S^1$. We would like to apply Eq. (2.31) to the $\mathcal{G}$-invariant function $\chi : \mathcal{A} \to \mathbb{C}$ given by

$$\chi(A) = \prod\nolimits_i \mathrm{Tr}_{\rho_i}\big(\mathrm{Hol}_{l_i}(A)\big) \exp(iS_{CS}(A)) \tag{2.33}$$

Before we can do this we need to extend the $\mathcal{G}_\Sigma$-invariant function $\chi^{qax} := \chi_{|\mathcal{A}^{qax}}$ on $\mathcal{A}^{qax}$ to a $\overline{\mathcal{G}_\Sigma}$-invariant function $\overline{\chi^{qax}}$ on $\overline{\mathcal{A}^{qax}}$. We do this by extending each of the $\mathcal{G}_\Sigma$-invariant functions

$$\mathrm{Tr}_{\rho_i}(\mathrm{Hol}_{l_i}) : \mathcal{A}^{qax} \ni A^q \mapsto \mathrm{Tr}_{\rho_i}\big(\mathrm{Hol}_{l_i}(A^q)\big) \in \mathbb{C}, \qquad i \leq m \tag{2.34}$$

$$S^{qax}_{CS} : \mathcal{A}^{qax} \ni A^q \mapsto S_{CS}(A^q) \in \mathbb{C} \tag{2.35}$$

separately to $\overline{\mathcal{G}_\Sigma}$-invariant functions $\overline{\mathrm{Tr}_{\rho_i}(\mathrm{Hol}_{l_i})}$ and $\overline{S^{qax}_{CS}}$ on $\overline{\mathcal{A}^{qax}}$ and then setting

$$\overline{\chi^{qax}}(A^q) := \prod\nolimits_i \overline{\mathrm{Tr}_{\rho_i}(\mathrm{Hol}_{l_i})}(A^q) \exp(i\overline{S^{qax}_{CS}}(A^q)) \tag{2.36}$$

for $A^q \in \overline{\mathcal{A}^{qax}}$. If σ_0 is not in the image of the loops l^j_Σ, which we will assume in the following, then $\mathrm{Hol}_{l_i}(A^q)$ makes sense for arbitrary[17] $A^q \in \overline{\mathcal{A}^{qax}}$ and setting $\overline{\mathrm{Tr}_{\rho_i}(\mathrm{Hol}_{l_i})}(A^q) := \mathrm{Tr}_{\rho_i}(\mathrm{Hol}_{l_i}(A^q))$ we obtain a $\overline{\mathcal{G}_\Sigma}$-invariant function on $\overline{\mathcal{A}^{qax}}$.

Before we write down an explicit expression for $\overline{S^{qax}_{CS}}$ we will first give a convenient explicit formula for the function S^{qax}_{CS} on $\mathcal{A}^{qax} = \mathcal{A}^\perp \oplus C^\infty(\Sigma, \mathfrak{g})dt$. We have

$$\begin{aligned} S^{qax}_{CS}(A^\perp + Bdt) &= S_{CS}(A^\perp + Bdt) \\ &= k\pi \int_M \big[\mathrm{Tr}(A^\perp \wedge dA^\perp) + 2\,\mathrm{Tr}(A^\perp \wedge Bdt \wedge A^\perp) + 2\,\mathrm{Tr}(A^\perp \wedge dB \wedge dt)\big] \end{aligned} \tag{2.37}$$

for all $A^\perp \in \mathcal{A}^\perp$, $B \in C^\infty(\Sigma, \mathfrak{g})$. As in [24, 26] we can rewrite the RHS of Eq. (2.37) using the identification

$$\mathcal{A}^\perp \cong C^\infty(S^1, \mathcal{A}_\Sigma) \tag{2.38}$$

where $C^\infty(S^1, \mathcal{A}_\Sigma)$ denotes the space of all functions $\alpha : S^1 \to \mathcal{A}_\Sigma$ which are "smooth" in the sense that for every smooth vector field X on Σ the function $\Sigma \times S^1 \ni (\sigma, t) \mapsto \alpha(t)(X_\sigma)$ is smooth.

Using this identification we have (cf. Proposition 5.2 in [24])

$$S^{qax}_{CS}(A^\perp + Bdt) = -k\pi \int_{S^1} dt \int_\Sigma \Big[\mathrm{Tr}\big(A^\perp(t) \wedge \big(\tfrac{\partial}{\partial t} + \mathrm{ad}(B)\big)A^\perp(t)\big) - 2\,\mathrm{Tr}\big(A^\perp(t) \wedge dB\big)\Big] \tag{2.39}$$

where $\frac{\partial}{\partial t} : C^\infty(S^1, \mathcal{A}_\Sigma) \to C^\infty(S^1, \mathcal{A}_\Sigma)$ is the obvious differential operator.

Let us now introduce $\overline{S^{qax}_{CS}} : \overline{\mathcal{A}^{qax}} \to \mathbb{C}$ by[18]

$$\begin{aligned} &\overline{S^{qax}_{CS}}(A^\perp + Bdt) \\ &\quad := -k\pi \Big\{ \lim_{\epsilon \to 0} \int_{S^1} dt \int_{\Sigma \backslash B_\epsilon(\sigma_0)} \Big[\mathrm{Tr}\big(A^\perp(t) \wedge \big(\tfrac{\partial}{\partial t} + \mathrm{ad}(B)\big)A^\perp(t)\big) - 2\,\mathrm{Tr}\big(dA^\perp(t) \cdot B\big)\Big]\Big\} \end{aligned} \tag{2.40}$$

---

[17] in fact this expression makes sense even for arbitrary $A^q \in \mathcal{A}^{qax}_{(\Sigma\backslash\{\sigma_0\})\times S^1}$ and defines a $\mathcal{G}_{\Sigma\backslash\{\sigma_0\}}$-invariant function $\overline{\mathrm{Tr}_{\rho_i}(\mathrm{Hol}_{l_i})}$ on all $\mathcal{A}^{qax}_{(\Sigma\backslash\{\sigma_0\})\times S^1}$

[18] here $B_\epsilon(\sigma_0)$ is the closed ϵ-ball around σ_0 w.r.t. to an arbitrary fixed auxiliary Riemannian metric on Σ

for all $A^{\perp} \in \overline{\mathcal{A}^{\perp}}$, $B \in \overline{C^{\infty}}(\Sigma, \mathfrak{g})$ for which the $\epsilon \to 0$-limit exists and by

$$\overline{S^{qax}_{CS}}(A^{\perp} + Bdt) := 0$$

otherwise. Observe that $\overline{S^{qax}_{CS}}$ is indeed an extension of S^{qax}_{CS} since in the special case where $A \in \mathcal{A}^{\perp}$ and $B \in C^{\infty}(\Sigma, \mathfrak{g})$ Stokes' Theorem implies that for every $t \in S^1$ we have[19]

$$\int_{\Sigma \backslash \{\sigma_0\}} \mathrm{Tr}\big(dA^{\perp}(t) \cdot B\big) = \int_{\Sigma} \mathrm{Tr}\big(dA^{\perp}(t) \cdot B\big) = \int_{\Sigma} \mathrm{Tr}\big(A^{\perp}(t) \wedge dB\big) \tag{2.41}$$

Moreover, $\overline{S^{qax}_{CS}}$ is indeed a $\overline{\mathcal{G}_{\Sigma}}$-invariant function, cf. Proposition 4.1 in [25]. We can therefore apply Eq. (2.31) to the functions χ and $\overline{\chi^{qax}}$ given by Eq. (2.33) and Eq. (2.36) with the choice Eq. (2.40) above and obtain

$$\begin{aligned}\mathrm{WLO}(L) \sim \sum\nolimits_{\mathrm{h} \in [\Sigma, G/T]} \int_{C^{\infty}(\Sigma, \mathfrak{t}_{reg})} \Big[\int_{\mathcal{A}^{\perp}} \prod\nolimits_i \overline{\mathrm{Tr}_{\rho_i}(\mathrm{Hol}_{l_i})}\big(A^{\perp} + A_{\mathrm{sg}}(\mathrm{h}) + Bdt\big) \\ \times \exp(i\overline{S^{qax}_{CS}}(A^{\perp} + A_{\mathrm{sg}}(\mathrm{h}) + Bdt)) DA^{\perp} \Big] \det\big(1_{\mathfrak{k}} - \exp(\mathrm{ad}(B))_{|\mathfrak{k}}\big) DB\end{aligned} \tag{2.42}$$

Here and in the following $\sim$ denotes equality up to a multiplicative constant independent of L.

Using the short notation $\mathrm{Tr}_{\rho_i}(\mathrm{Hol}_{l_i})$ instead of $\overline{\mathrm{Tr}_{\rho_i}(\mathrm{Hol}_{l_i})}$ and taken into account that

$$\overline{S^{qax}_{CS}}(A^{\perp} + A_{\mathrm{sg}}(\mathrm{h}) + Bdt) = S_{CS}(A^{\perp} + Bdt) + 2\pi k \int_{\Sigma \backslash \{\sigma_0\}} \mathrm{Tr}\big(dA_{\mathrm{sg}}(\mathrm{h}) \cdot B\big) \tag{2.43}$$

we can rewrite Eq. (2.42) in the following form:

$$\begin{aligned}&\mathrm{WLO}(L) \\ &\sim \sum\nolimits_{\mathrm{h} \in [\Sigma, G/T]} \int_{C^{\infty}(\Sigma, \mathfrak{t}_{reg})} \Big[\int_{\mathcal{A}^{\perp}} \prod\nolimits_i \mathrm{Tr}_{\rho_i}\big(\mathrm{Hol}_{l_i}(A^{\perp} + A_{\mathrm{sg}}(\mathrm{h}) + Bdt)\big) \exp(iS_{CS}(A^{\perp} + Bdt)) DA^{\perp} \Big] \\ &\qquad \times \exp\big(2\pi i k \int_{\Sigma \backslash \{\sigma_0\}} \mathrm{Tr}\big(dA_{\mathrm{sg}}(\mathrm{h}) \cdot B\big)\big) \det\big(1_{\mathfrak{k}} - \exp(\mathrm{ad}(B))_{|\mathfrak{k}}\big) DB\end{aligned} \tag{2.44}$$

Remark 2.6 One could think that it is possible to simplify the RHS of Eq. (2.42) by making, for every fixed $\mathrm{h} \in [\Sigma, G/T]$, the informal change of variable $A^{\perp} + A_{\mathrm{sg}}(\mathrm{h}) \to A^{\perp}$. However, this change of variable is not admissible, cf. Sec. 4.2 in [25]. In fact, it turns out that the explicit evaluation of the simplified expression obtained in this way leads to "incorrect" values for WLO(L).

On the other hand, it might be possible to justify a similar change of variable using a slightly different procedure. In order to do so let us consider the extension

$$V := \mathcal{A}^{qax}(T) \oplus \big(\oplus_{\mathrm{h} \in [\Sigma, G/T]} \mathbb{R} \cdot A_{\mathrm{sg}}(\mathrm{h})\big)$$

of $\mathcal{A}^{qax}(T)$ and introduce the extension $S^{V}_{CS} : V \to \mathbb{C}$ of $S^{qax}_{CS} : \mathcal{A}^{qax}(T) \to \mathbb{C}$ by[20]

$$\begin{aligned}S^{V}_{CS}(A^{\perp} + A_{\mathrm{sg}}(\mathrm{h}) + Bdt) := -k\pi \int_{S^1} dt \int_{\Sigma \backslash \{\sigma_0\}} \Big[\mathrm{Tr}\big((A^{\perp}(t) + A_{\mathrm{sg}}(\mathrm{h})) \wedge \big(\tfrac{\partial}{\partial t} + \mathrm{ad}(B)\big)(A^{\perp}(t) + A_{\mathrm{sg}}(\mathrm{h}))\big) \\ - 2\,\mathrm{Tr}\big((A^{\perp}(t) + A_{\mathrm{sg}}(\mathrm{h})) \wedge dB\big) \Big]\end{aligned}$$

---

[19] here and above $dA^{\perp}(t)$ is a short notation for $d(A^{\perp}(t))$, i.e. "d" is the differential of Σ

[20] here $\int_{\Sigma \backslash \{\sigma_0\}} \cdots$ is a short notation for $\lim_{\epsilon \to 0} \int_{\Sigma \backslash B_{\epsilon}(\sigma_0)} \cdots$. Observe that this $\epsilon \to 0$-limit will always exist (in contrast to the situation in Eq. (2.40) above)

for all $A^\perp \in \mathcal{A}^\perp$, $B \in C^\infty(\Sigma, \mathfrak{t})$, and $\mathrm{h} \in [\Sigma, G/T]$. Observe that we have

$$\overline{S^{qax}_{CS}}(A^\perp + A_{\mathrm{sg}}(\mathrm{h}) + Bdt) = S^V_{CS}(A^\perp + A_{\mathrm{sg}}(\mathrm{h}) + Bdt) + 2\pi k \operatorname{Tr}\big(n(\mathrm{h}) \cdot B(\sigma_0)\big) \tag{2.45}$$

where $n(\mathrm{h}) \in \mathfrak{t}$ is given as in Remark 2.5 above. If we replace the expression $\overline{S^{qax}_{CS}}(A^\perp + A_{\mathrm{sg}}(\mathrm{h}) + Bdt)$ in Eq. (2.42) by the RHS of Eq. (2.45), perform for each fixed $\mathrm{h} \in [\Sigma, G/T]$ the change of variable $A^\perp + A_{\mathrm{sg}}(\mathrm{h}) \to A^\perp$ and use the relation $S^V_{CS}(A^\perp + Bdt) = S_{CS}(A^\perp + Bdt)$ for $A^\perp \in \mathcal{A}^\perp$, $B \in C^\infty(\Sigma, \mathfrak{t})$ we obtain

$$\begin{aligned}\mathrm{WLO}(L) \sim \sum\nolimits_{\mathrm{h} \in [\Sigma, G/T]} \int_{C^\infty(\Sigma, \mathfrak{t}_{reg})} \Big[\int_{\mathcal{A}^\perp} \prod\nolimits_i \operatorname{Tr}_{\rho_i}\big(\mathrm{Hol}_{l_i}(A^\perp + Bdt)\big) \exp(iS_{CS}(A^\perp + Bdt)) DA^\perp \Big] \\ \times \exp\big(2\pi i k \operatorname{Tr}(n(\mathrm{h}) \cdot B(\sigma_0))\big) \det\big(1_{\mathfrak{k}} - \exp(\mathrm{ad}(B))_{|\mathfrak{k}}\big) DB \end{aligned} \tag{2.46}$$

It turns out that the evaluation of the RHS of Eq. (2.46) (either using the continuum computations in [26, 18] or by proceeding as in the present paper and in [28]) *does* lead to the correct result for WLO(L), cf. Remark 2.2 in [28]. However, since this issue is not totally understood at the present time we prefer to work with Eq. (2.44) above rather than with Eq. (2.46) during the main part of the present paper (and [28]).

2.3.2 Rewriting $S_{CS}(A^\perp + Bdt)$

Let us rewrite $S_{CS}(A^\perp + Bdt)$ yet another time. In order to do so we fix an auxiliary Riemannian metric $\mathbf{g}$ on Σ. Then Eq. (2.39) can be rewritten as (cf.[21] Sec. 3.3 in [26])

$$S_{CS}(A^\perp + Bdt) = k\pi \Big[\ll A^\perp, \star\big(\tfrac{\partial}{\partial t} + \mathrm{ad}(B)\big) A^\perp \gg_{\mathcal{A}^\perp} - 2 \ll A^\perp, \star dB \gg_{\mathcal{A}^\perp} \Big] \tag{2.47}$$

where $\star$ denotes both the Hodge star operator on $\mathcal{A}_\Sigma$ induced by $\mathbf{g}$ and the linear isomorphism on $\mathcal{A}^\perp \cong C^\infty(S^1, \mathcal{A}_\Sigma)$ given by $(\star A^\perp)(t) = \star(A^\perp(t))$ for all $t \in S^1$. Above $\ll \cdot, \cdot \gg_{\mathcal{A}^\perp}$ is the scalar product on $\mathcal{A}^\perp \cong C^\infty(S^1, \mathcal{A}_\Sigma)$ given by

$$\ll A^\perp_1, A^\perp_2 \gg_{\mathcal{A}^\perp} = \int_{S^1} dt \int_\Sigma (A^\perp_1(t), A^\perp_2(t))_{\mathcal{A}_\Sigma} d\mu_{\mathbf{g}} \tag{2.48}$$

where $d\mu_{\mathbf{g}}$ is the volume measure on Σ associated to $\mathbf{g}$ and $(\cdot, \cdot)_{\mathcal{A}_\Sigma} : \mathcal{A}_\Sigma \times \mathcal{A}_\Sigma \to C^\infty(\Sigma, \mathbb{R})$ is the obvious bilinear map induced by $\mathbf{g}$ and $\langle \cdot, \cdot \rangle_{\mathfrak{g}}$.

We remark that even though Eq. (2.47) is less natural than Eq. (2.39) above (since it depends on the non-canonically chosen Riemannian metric $\mathbf{g}$) it will be more convenient for our purposes, cf. the paragraph after Remark 2.7 below and also Sec. 5.2 below where a discrete analogue of $S_{CS}(A^\perp + Bdt)$ is introduced.

2.3.3 The decomposition $\mathcal{A}^\perp = \check{\mathcal{A}}^\perp \oplus \mathcal{A}^\perp_c$

Clearly, the operator $\frac{\partial}{\partial t} + \mathrm{ad}(B) : C^\infty(S^1, \mathcal{A}_\Sigma) \to C^\infty(S^1, \mathcal{A}_\Sigma)$ is not injective. For $B \in C^\infty(\Sigma, \mathfrak{t}_{reg})$ the kernel of this operator is given by

$$\mathcal{A}^\perp_c := \{A^\perp \in C^\infty(S^1, \mathcal{A}_\Sigma) \mid A^\perp \text{ is constant and } \mathcal{A}_{\Sigma,\mathfrak{t}}\text{-valued}\} \cong \mathcal{A}_{\Sigma,\mathfrak{t}} \tag{2.49}$$

For making rigorous sense of the RHS of Eq. (2.44) it is useful to work with a decomposition $\mathcal{A}^\perp = \mathcal{C} \oplus \mathcal{A}^\perp_c$ for a suitably chosen linear subspace $\mathcal{C}$ of $\mathcal{A}^\perp$ and to incorporate this decomposition into Eq. (2.44). For technical reasons we worked in [24, 25, 26, 27] with the choice $\mathcal{C} := \hat{\mathcal{A}}^\perp$ where

$$\hat{\mathcal{A}}^\perp := \{A^\perp \in C^\infty(S^1, \mathcal{A}_\Sigma) \mid A^\perp(t_0) \in \mathcal{A}_{\Sigma,\mathfrak{k}}\} \tag{2.50}$$

---

[21] we remark that in [26] we use a different sign convention for the $\star$ operator

where t_0 is an arbitrary but fixed point in S^1. It would also have been possible to work with $\mathcal{C} := \check{\mathcal{A}}^{\perp}$ where

$$\check{\mathcal{A}}^{\perp} := \{A^{\perp} \in C^{\infty}(S^1, \mathcal{A}_{\Sigma}) \mid \int_{S^1} A^{\perp}(t) dt \in \mathcal{A}_{\Sigma,\mathfrak{k}}\} \tag{2.51}$$

In fact, the latter choice is in some sense more natural and has the important advantage of possessing a "good" discrete analogue, cf. the second of the two equations in (5.25) below. In the present paper we will therefore work with the latter choice. Taking into account that

$$S_{CS}(\check{A}^{\perp} + A^{\perp}_c + Bdt) = S_{CS}(\check{A}^{\perp} + Bdt) + S_{CS}(A^{\perp}_c + Bdt) \tag{2.52}$$

for $\check{A}^{\perp} \in \check{\mathcal{A}}^{\perp}$, $A^{\perp}_c \in \mathcal{A}^{\perp}_c$, and $B \in \mathcal{B}$ we arrive, after incorporating the decomposition $\mathcal{A}^{\perp} = \check{\mathcal{A}}^{\perp} \oplus \mathcal{A}^{\perp}_c$ into Eq. (2.44), at

$$\begin{aligned}
\mathrm{WLO}(L) \sim \textstyle\sum_{\mathrm{h} \in [\Sigma, G/T]} \int_{\mathcal{A}^{\perp}_c \times \mathcal{B}} \Big\{ & 1_{C^{\infty}(\Sigma, \mathfrak{t}_{reg})}(B) \, \mathrm{Det}_{FP}(B) \\
& \times \Big[\int_{\check{\mathcal{A}}^{\perp}} \textstyle\prod_i \mathrm{Tr}_{\rho_i}\big(\mathrm{Hol}_{l_i}(\check{A}^{\perp} + A^{\perp}_c, B; \mathrm{h})\big) \exp(iS_{CS}(\check{A}^{\perp}, B)) D\check{A}^{\perp} \Big] \\
& \times \exp\big(2\pi i k \int_{\Sigma \backslash \{\sigma_0\}} \mathrm{Tr}\big(dA_{\mathrm{sg}}(\mathrm{h}) \cdot B\big)\big) \Big\} \exp(iS_{CS}(A^{\perp}_c, B))(DA^{\perp}_c \otimes DB)
\end{aligned} \tag{2.53}$$

where, as a preparation for Sec. 5, we have introduced the short notation

$$S_{CS}(A^{\perp}, B) := S_{CS}(A^{\perp} + Bdt) \tag{2.54a}$$

$$\mathrm{Det}_{FP}(B) := \det\big(1_{\mathfrak{k}} - \exp(\mathrm{ad}(B))_{|\mathfrak{k}}\big) \tag{2.54b}$$

$$\mathrm{Hol}_l(A^{\perp}, B; \mathrm{h}) := \mathrm{Hol}_l(A^{\perp} + A_{\mathrm{sg}}(\mathrm{h}) + Bdt) \tag{2.54c}$$

(for $l = l_i$ and $A^{\perp} = \check{A}^{\perp} + A^{\perp}_c$) and where we have rewritten the integral $\int_{C^{\infty}(\Sigma, \mathfrak{t}_{reg})} \cdots$ in the form $\int_{\mathcal{B}} 1_{C^{\infty}(\Sigma, \mathfrak{t}_{reg})}(B) \cdots$.

Remark 2.7 By generalizing the heuristic arguments above in an obvious way Eq. (2.53) can also be "derived" for general simply-connected compact $G = \prod_{i=1}^{r} G_i$, cf. Remark 2.2, provided that we replace the first equation in (2.54) by $S_{CS}(A^{\perp}, B) := S_{CS}(M, G, (k_i)_i)(A^{\perp}, B) := S_{CS}(M, \prod_i G_i, (k_i)_i)(A^{\perp} + Bdt)$ where $(G_i)_{i \leq r}$ and $(k_i)_{i \leq r}$ are as in Remark 2.2.

From the explicit formula Eq. (2.47) for $S_{CS}(A^{\perp} + Bdt)$ it follows immediately that both (heuristic) complex measures $\exp(iS_{CS}(\check{A}^{\perp}, B))D\check{A}^{\perp}$ and $\exp(iS_{CS}(A^{\perp}_c, B))(DA^{\perp}_c \otimes DB)$ appearing above are of "Gaussian type". This considerably increases the chances of making rigorous sense of the RHS of Eq. (2.53).

In fact, in [24, 26, 27] we have already sketched how this works in the framework of white noise analysis (see part F of the Appendix below for some comments on this approach). In [24, 26, 27, 18] we also demonstrated that in the special case where the link L has no double points and "horizontal" framing is used (cf. Sec. 5.2 in [26]) the rigorous realization of $\mathrm{WLO}(L)$ can be evaluated explicitly and that

$$\mathrm{WLO}(L) \sim |L| \tag{2.55}$$

holds where $|L|$ is Turaev's shadow invariant for the group G (cf. part B of the Appendix in [28]) and where $\sim$ denotes equality up to a multiplicative constant independent of L.

In the present paper and its sequel [28] we will develop an alternative "discretization" approach for making rigorous sense of the RHS of Eq. (2.53), cf. Theorem 5.3 in [28] which is a rigorous and elementary version of Eq. (2.55) for the same class of links as in [24, 26, 27].

3 Oscillatory Gauss-type measures on Euclidean spaces

Let us fix a Euclidean vector space $(V, \langle \cdot, \cdot \rangle)$ and set $d := \dim(V)$.

3.1 Basic Definitions

Definition 3.1 *An "oscillatory Gauss-type measure" on* $(V, \langle \cdot, \cdot \rangle)$ *is a complex Borel measure* $d\mu$ *on* V *of the form*

$$d\mu(x) = \tfrac{1}{Z} e^{-\frac{i}{2}\langle x-m, S(x-m)\rangle} dx \tag{3.1}$$

with $Z \in \mathbb{C}\backslash\{0\}$, $m \in V$, *and where* S *is a symmetric endomorphism of* V *and* dx *the normalized*[22] *Lebesgue measure on* V. *Note that* Z, m *and* S *are uniquely determined by* $d\mu$ *so we can use the notation* Z_μ, m_μ *and* S_μ *in order to denote these objects.*

i) We call $d\mu$ *"centered"iff* $m = 0$.

ii) We call $d\mu$ *"degenerate" iff* S *is not invertible*

iii) We call $d\mu$ *"normalized" iff* $Z = \frac{(2\pi)^{d/2}}{\det^{\frac{1}{2}}(iS')}$ *where* $S' := S_{|\ker(S)^\perp}$. *(See Example 3.4 below for the definition of* $\det^{\frac{1}{2}}(iS)$ *and a motivation for the term "normalized").*

Definition 3.2 *Let* $d\mu$ *be an oscillatory Gauss-type measure on* $(V, \langle \cdot, \cdot \rangle)$. *A (Borel) measurable function* $f : V \to \mathbb{C}$ *will be called improperly integrable w.r.t.* $d\mu$ *iff*[23]

$$\int_\sim f d\mu := \int_\sim f(x) d\mu(x) := \lim_{\epsilon \to 0} (\tfrac{\epsilon}{\pi})^{n/2} \int f(x) e^{-\epsilon |x|^2} d\mu(x) \tag{3.2}$$

exists. Here we have set $n := \dim(\ker(S_\mu))$. *Note that if* $d\mu$ *is non-degenerate we have* $n = 0$ *so the factor* $(\frac{\epsilon}{\pi})^{n/2}$ *is then trivial.*

Most of the time we will consider non-degenerate oscillatory Gauss-type measures, the exception being Proposition 3.12 below.

Using a simple analytic continuation argument and the corresponding explicit formulas for Gaussian probability measures we can easily prove the existence of $\int_\sim f \, d\mu$ and compute the corresponding value explicitly for a large class of functions f. Let us illustrate this by looking at some simple examples:

Example 3.3 *Consider the special case where* $V = \mathbb{R}$, *where* $\langle \cdot, \cdot \rangle$ *is the scalar product given by* $\langle x, y \rangle = x \cdot y$, *and where* $d\mu(x) = \exp(i\langle x, x\rangle)dx = \exp(ix^2)dx$. *Then the improper integrals*

$$\int_\sim 1 \ d\mu(x), \quad \int_\sim x \ d\mu(x), \quad \int_\sim x^2 \ d\mu(x), \quad \int_\sim e^{cx} \ d\mu(x), \ c \in \mathbb{C}$$

exist and are given explicitly by

- $\int_\sim 1 \ d\mu(x) = \sqrt{i\pi}$
- $\int_\sim x \ d\mu(x) = 0$
- $\int_\sim x^2 \ d\mu(x) = \frac{i}{2} \sqrt{i\pi}$

---

[22] i.e. unit hyper-cubes have volume 1 w.r.t. dx

[23] Observe that $\int_{\ker(S_\mu)} e^{-\epsilon\|x\|^2} d\mu(x) = (\frac{\epsilon}{\pi})^{-n/2}$. In particular, the factor $(\frac{\epsilon}{\pi})^{n/2}$ in Eq. (3.2) above ensures that also for degenerate oscillatory Gauss-type measure the improper integrals $\int_\sim 1 \ d\mu$ exists, cf. Example 3.4 below

- $\int_{\sim} e^{cx} \ d\mu(x) = e^{i\frac{c^2}{4}} \ \sqrt{i\pi}$

where $\sqrt{\cdot} : \mathbb{C}\backslash(-\infty, 0) \to \mathbb{C}$ *denotes the standard square root.*

In order to show the existence (and to compute the explicit value) of $\int_{\sim} 1 \ d\mu(x)$ *we consider the analytic function* $F : \{z \mid Re(z) > 0\} \to \mathbb{C}$ *given by* $F(z) := \int \exp(-zx^2)dx$*. According to a well-known formula we have* $F(a) = \sqrt{\pi/a}$ *for all* $a \in (0, \infty)$*. The obvious uniqueness argument for analytic functions now implies that* $F(z) = \sqrt{\pi/z}$ *for all* $z \in \mathbb{C}$ *with* $Re(z) > 0$*. Thus* $\int_{\sim} 1 \ d\mu = \lim_{\epsilon \to 0} F(\epsilon - i) = \lim_{\epsilon \to 0} \sqrt{\pi/(\epsilon - i)} = \sqrt{i\pi} = \sqrt{\pi} \ e^{\frac{\pi}{4}i}$

The other three integrals can be dealt with in a similar way.

In the next example $(V, \langle \cdot, \cdot \rangle)$ is again an arbitrary Euclidean space.

Example 3.4 *Let* $d\mu$ *be a non-degenerate oscillatory Gauss-type measure on* $(V, \langle \cdot, \cdot \rangle)$ *with* S*,* m*, and* Z *given as in Eq.* (3.1)

i) We have[24]

$$\int_{\sim} 1 \ d\mu = \frac{1}{Z} \frac{(2\pi)^{d/2}}{\det^{\frac{1}{2}}(iS)} \tag{3.3}$$

where we have set $\det^{\frac{1}{2}}(iS) := \prod_k \sqrt{i\lambda_k} = e^{\frac{\pi i}{4} \sum_k \operatorname{sgn}(\lambda_k)} (\prod_k |\lambda_k|^{1/2})$ *where* $(\lambda_k)_k$ *are the (real) eigenvalues of the symmetric matrix* S*. In particular,* $d\mu$ *is normalized in the sense of Definition 3.1 above iff* $\int_{\sim} 1 \ d\mu = 1$*.*

ii) In the special case when $d\mu$ *is normalized we have for all* $v, w \in V$

$$\int_{\sim} \langle v, x \rangle \ d\mu(x) = \langle v, m \rangle, \qquad \int_{\sim} \langle v, x \rangle \langle w, x \rangle \ d\mu(x) = \tfrac{1}{i} \langle v, S^{-1} w \rangle + \langle m, m \rangle \tag{3.4}$$

We will not try to identify the largest possible class of functions f for which $\int_{\sim} f \ d\mu$ exists. For our purposes the function algebra $\mathcal{P}_{exp}(V)$ defined in the next definition will be sufficient.

Definition 3.5 *i) Let* W *be a finite-dimensional associative* $\mathbb{R}$*-algebra (with the standard topology). By* $\mathcal{P}_{exp}(V, W)$ *we will denote the subalgebra of* $\operatorname{Map}(V, W)$ *which is generated by the affine maps* $\varphi : V \to W$ *and their "exponentials"* $\exp_W \circ \varphi$*. Here* $\exp_W : W \to W$ *denotes the exponential map of* W*.*

ii) By $\mathcal{P}_{exp}(V)$ *we denote the subalgebra of* $\operatorname{Map}(V, \mathbb{C})$ *which is generated by the functions of the form* $\theta \circ f$ *with* $f \in \mathcal{P}_{exp}(V, W)$ *and* $\theta \in \operatorname{Hom}_{\mathbb{R}}(W, \mathbb{C})$ *where* W *is any*[25] *finite-dimensional associative* $\mathbb{R}$*-algebra.*

Using analytic continuation arguments, some explicit formulas for Gaussian probability measures, and suitable growth estimates one can prove the following result (which is easy to believe):

Proposition 3.6 *Let* $d\mu$ *be a non-degenerate oscillatory Gauss-type measure on* $(V, \langle \cdot, \cdot \rangle)$*. Then for every* $f \in \mathcal{P}_{exp}(V)$ *the improper integral* $\int_{\sim} f \ d\mu \in \mathbb{C}$ *exists.*

---

[24] we remark that if $d\mu$ is degenerate then an analogous statement will hold with S replaced by $S' := S_{| \ker(S)^{\perp}}$

[25] note that we do not keep W fixed here

3.2 Three propositions

Let us fix for a while a normalized non-degenerate oscillatory Gauss-type measure $d\mu$ on $(V, \langle\cdot,\cdot\rangle)$ and introduce the notation

$$\mathbb{E}_{\sim}[X] := \int_{\sim} X d\mu \in \mathbb{C} \tag{3.5a}$$

$$\mathrm{cov}_{\sim}(X, X') := \mathbb{E}_{\sim}[XX'] - \mathbb{E}_{\sim}[X]\,\mathbb{E}_{\sim}[X'] \in \mathbb{C} \tag{3.5b}$$

for maps $X, X' \in \mathcal{P}_{exp}(V)$ (in analogy to the case of (Gaussian or non-Gaussian) probability measures on V).

Observation 3.7 *Let $d\mu$ be as above and let $X_1, X_2, \ldots, X_n$ be a sequence of affine maps $V \to \mathbb{R}$.*

i) If $\mathbb{E}_{\sim}[X_i] = 0$ for every $i \leq n$ then

$$\mathbb{E}_{\sim}[\prod_j X_j] = \begin{cases} \frac{1}{(n/2)!2^{n/2}} \sum_{\sigma \in S_n} \prod_{i=1}^{n/2} \mathrm{cov}_{\sim}(X_{\sigma(2i-1)}, X_{\sigma(2i)}) & \textit{if } n \textit{ is even} \\ 0 & \textit{if } n \textit{ is odd} \end{cases} \tag{3.6}$$

ii) If $\mathrm{cov}_{\sim}(X_i, X_j) = 0$ for $i, j \leq n$ with $i \neq j$ we have

$$\mathbb{E}_{\sim}[\prod_j X_j] = \prod_j \mathbb{E}_{\sim}[X_j] \tag{3.7}$$

Here Eq. (3.6) follows from the analogous formula for the moments of a Gaussian probability measure and a suitable analytic continuation argument. Clearly, in the special case where $\mathrm{cov}_{\sim}(X_i, X_j) = 0$ for $i, j \leq n$ with $i \neq j$ Eq. (3.6) reduces to $\mathbb{E}_{\sim}[\prod_j X_j] = 0$. By applying the latter equation to the subsequences of the sequence $X'_1, X'_2, \ldots, X'_n$ given by $X'_j := X_j - \mathbb{E}[X_j]$ we arrive at Eq. (3.7).

Let us now consider the case where $Y : V \to \mathbb{R}$ is an affine map with[26] $\mathrm{cov}_{\sim}(Y, Y) = 0$ and let us consider the trivial sequence $(X_i)_{i=1}^n$ where $X_i = Y$ for each $i \leq n$. Since $\mathrm{cov}(Y, Y)_{\sim} = 0$ we trivially have $\mathrm{cov}_{\sim}(X_i, X_j) = 0$ for $i \neq j$ (and even for $i = j$) so according to Eq. (3.7) we have $\mathbb{E}_{\sim}[Y^n] = \mathbb{E}_{\sim}[Y]^n$, from which we conclude, for example, that

$$\mathbb{E}_{\sim}[\exp(Y)] = \mathbb{E}_{\sim}[\sum_n \tfrac{Y^n}{n!}] \overset{(*)}{=} \sum_n \mathbb{E}_{\sim}[\tfrac{Y^n}{n!}] = \sum_n \frac{\mathbb{E}_{\sim}[Y]^n}{n!} = \exp(\mathbb{E}_{\sim}[Y]) \tag{3.8}$$

In order to see that step $(*)$ above holds observe that for every fixed $\epsilon > 0$ we have

$$\int \sum_n \left| \tfrac{Y(x)^n}{n!} e^{-\epsilon|x|^2} \tfrac{1}{Z} e^{-\frac{i}{2}\langle x-m, S(x-m)\rangle} \right| dx \leq \int \sum_n \tfrac{|Y(x)|^n}{n!} \tfrac{1}{Z} e^{-\epsilon|x|^2} dx < \infty$$

and we can therefore conclude from the dominated convergence theorem[27] that

$$\int \sum_n \tfrac{Y(x)^n}{n!} e^{-\epsilon|x|^2} d\mu(x) = \sum_n \int \tfrac{Y(x)^n}{n!} e^{-\epsilon|x|^2} d\mu(x)$$

for each $\epsilon > 0$. Accordingly, in order to prove step $(*)$ it is enough to prove that $\lim_{\epsilon \to 0} \sum_n \frac{I(n,\epsilon)}{n!} = \sum_n \lim_{\epsilon \to 0} \frac{I(n,\epsilon)}{n!}$ where we have set $I(n, \epsilon) := \int Y(x)^n e^{-\epsilon|x|^2} d\mu(x)$. The latter claim can easily be proven by computing the integrals $I(n, \epsilon)$ explicitly.

---

[26] Note that the condition $\mathrm{cov}_{\sim}(Y, Y) = 0$ does not imply that Y is a constant map on V. This is in sharp contrast to the situation for (Gaussian or non-Gaussian) probability measures where the relation $\mathrm{cov}(Y, Y) = 0$ always implies $Y = \mathbb{E}[Y]$ $d\mu$-a.s.

[27] applied to the positive measure dx "appearing" in $d\mu$

More generally, we obtain[28] for every $\Phi \in \mathcal{P}_{exp}(\mathbb{R})$

$$\mathbb{E}_{\sim}[\Phi(Y)] = \Phi(\mathbb{E}_{\sim}[Y]) \tag{3.9}$$

since every such Φ is necessarily entire analytic and the coefficients $(c_n)_n$, given by $\Phi(x) = \sum_{n=0}^{\infty} c_n x^n$ for all $x \in \mathbb{R}$, have the property that $c_n \overset{n\to\infty}{\longrightarrow} 0$ rapidly enough so that

1. we can again apply the dominated convergence theorem in a similar way as above and prove that the two limit procedures $\int \cdots dx$ and $\sum_n$ can be interchanged

2. we can prove again that the two limit procedures $\lim_{\epsilon\to 0}$ and $\sum_n$ can be interchanged

Finally, we can generalize Eq. (3.9) to the case where we have a $\Phi \in \mathcal{P}_{exp}(\mathbb{R}^n)$, $n \in \mathbb{N}$, and where $(Y_k)_{k\leq n}$ is a sequence of affine maps $V \to \mathbb{R}$ such that $\mathrm{cov}_{\sim}(Y_i, Y_j) = 0$ holds for all $i, j \leq n$. We then arrive at the following result, which will be the key argument in Sec. 5.4.1 in [28]. In order to make the application of Proposition 3.8 in Sec. 5.4.1 in [28] more transparent we avoid the use of the notation $\mathbb{E}_{\sim}[\cdot]$ and $\mathrm{cov}_{\sim}(\cdot,\cdot)$ from now on.

Proposition 3.8 *Let $d\mu$ be a normalized non-degenerate oscillatory Gauss-type measure on $(V, \langle\cdot,\cdot\rangle)$ and let $(Y_k)_{k\leq n}$, $n \in \mathbb{N}$, be a sequence of affine maps $V \to \mathbb{R}$ such that*

$$\int_{\sim} Y_i Y_j d\mu = \Big(\int_{\sim} Y_i d\mu\Big)\Big(\int_{\sim} Y_j d\mu\Big) \tag{3.10}$$

holds for all $i, j \leq n$. Then for every $\Phi \in \mathcal{P}_{exp}(\mathbb{R}^n)$ we have

$$\int_{\sim} \Phi((Y_k)_k) d\mu = \Phi\Big(\Big(\int_{\sim} Y_k d\mu\Big)_k\Big) \tag{3.11}$$

Remark 3.9 In Sec. 5.4.1 in [28] we will actually apply a reformulation of Proposition 3.8 where the sequence $(Y_k)_{k\leq n}$, $n \in \mathbb{N}$, is replaced by a family $(Y_k^{i,a})_{k\leq n, i\leq m, a\leq D}$ of affine maps fulfilling the obvious analogue of Eq. (3.10) above and the function $\Phi \in \mathcal{P}_{exp}(\mathbb{R}^n)$ is replaced by a function $\Phi \in \mathcal{P}_{exp}(\mathbb{R}^{m\times n\times D})$.

Example 3.10 *Consider the (non-degenerate normalized centered) oscillatory Gauss-type measure $d\mu(x) := \frac{1}{2\pi}\exp(i\langle x_1, x_2\rangle)dx_1 dx_2$ on $V = \mathbb{R}^2$. For every $f \in \mathcal{P}_{exp}(\mathbb{R})$ we have*

$$\int_{\sim} f(x_1) d\mu(x) = f(0) \tag{3.12}$$

This follows by applying Proposition 3.8 with $\Phi = f$ and $Y_1(x) := x_1$. Observe that Eq. (3.10) *is indeed fulfilled since according to Example 3.4 we have $\int_{\sim} Y_1 d\mu = \int_{\sim}\langle x, e_1\rangle d\mu(x) = 0$ and $\int_{\sim} Y_1 Y_1 d\mu = \frac{1}{i}\langle e_1, S_\mu^{-1} e_1\rangle = 0$ where we have set $e_1 := \begin{pmatrix} 1 \\ 0 \end{pmatrix}$ and used that $S_\mu = -\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$.*

Using a different argument one can easily prove that Eq. (3.12) holds for arbitrary continuous bounded functions f. Moreover, we can include an additional "exponential factor", and we can in fact consider more general oscillatory Gauss-type measures $d\mu$, cf. Proposition 3.12 below, which will play a key role in Sec. 5.4.2 in [28].

As a preparation for Proposition 3.12 let us first consider the special case where the oscillatory Gauss-type measure $d\mu$ is non-degenerate:

---

[28] Observe that $\Phi(Y) \in \mathcal{P}_{exp}(V)$ so the existence of the LHS of Eq. (3.9) is guaranteed by Proposition 3.6

Proposition 3.11 *Assume that $V = V_1 \oplus V_2$ where V_1 and V_2 are two subspaces of V which are orthogonal to each other. For each $j = 1, 2$ we denote the V_j-component of $x \in V$ by x_j. Moreover, let $d\mu$ be a (non-degenerate centered) normalized oscillatory Gauss-type measure on $(V, \langle \cdot, \cdot \rangle)$ of the form $d\mu(x) = \frac{1}{Z}\exp(i\langle x_2, Mx_1 \rangle)dx$ for some linear isomorphism $M : V_1 \to V_2$. Then for every bounded continuous function $f : V_1 \to \mathbb{C}$ and every fixed $v \in V_2$ we have*

$$\int_\sim f(x_1)\exp(i\langle x_2, v\rangle)d\mu(x) = f(-M^{-1}v) \tag{3.13}$$

(In particular, the LHS of Eq. (3.13) *exists; note that the present situation is not covered by Proposition 3.6 above).*

Proof. Let dx_1 resp. dx_2 be the normalized Lebesgue measure on V_1 resp. V_2 (equipped with the scalar product induced by the one on V). We have

$$\begin{aligned}
&\int_\sim f(x_1)\exp(i\langle x_2, v\rangle)d\mu(x) \\
&= \frac{1}{Z}\lim_{\epsilon \to 0}\int_{V_1}\int_{V_2} e^{-\epsilon(|x_1|^2+|x_2|^2)} f(x_1)\exp(i\langle x_2, v + Mx_1\rangle)dx_2 dx_1 \\
&= \frac{(2\pi)^{d_2}}{Z}\lim_{\epsilon \to 0}\int_{V_1} e^{-\epsilon|x_1|^2} f(x_1)\delta_\epsilon(v + Mx_1)dx_1
\end{aligned} \tag{3.14}$$

where $d_2 := \dim(V_2) = d/2$ and where $\delta_\epsilon : V_2 \to \mathbb{R}$ is given by

$$\delta_\epsilon(w) := \tfrac{1}{(2\pi)^{d_2}}\int_{V_2} e^{-\epsilon|x_2|^2}\exp(i\langle x_2, w\rangle)dx_2 = \tfrac{1}{(2\pi)^{d_2}}\left(\tfrac{\pi}{\epsilon}\right)^{d_2/2} e^{-\frac{1}{2}\frac{|w|^2}{2\epsilon}} = \tfrac{1}{(4\epsilon\pi)^{d_2/2}} e^{-\frac{|w|^2}{4\epsilon}} \tag{3.15}$$

for all $w \in V_2$. Let us now fix an (arbitrary) isometry $\psi : V_2 \to V_1$. Clearly, the pushforward $(\psi)_* dx_2$ of dx_2 coincides with dx_1 so we have

$$\int_{V_1} e^{-\epsilon|x_1|^2} f(x_1)\delta_\epsilon(v + Mx_1)dx_1 = \int_{V_2} e^{-\epsilon|\psi x_2|^2} f(\psi x_2)\delta_\epsilon(v + M\psi x_2)dx_2 \tag{3.16}$$

Making the change of variable $M\psi x_2 + v \to y_2$ on the RHS of Eq. (3.16) we obtain

$$\int_{V_1} e^{-\epsilon|x_1|^2} f(x_1)\delta_\epsilon(v + Mx_1)dx_1 = |\det(M\psi)|^{-1}\int_{V_2} e^{-\epsilon|M^{-1}(y_2-v)|^2} f(M^{-1}(y_2 - v))\delta_\epsilon(y_2)dy_2 \tag{3.17}$$

where dy_2 is the normalized Lebesgue measure on V_2. Since $(\delta_\epsilon)_{\epsilon>0}$ is an "approximation to the identity" (i.e. converges weakly to the Dirac distribution δ_0) it is therefore clear[29] that

$$\lim_{\epsilon \to 0}\int_{V_1} e^{-\epsilon|x_1|^2} f(x_1)\delta_\epsilon(v + Mx_1)dx_1 = f(-M^{-1}v) \cdot |\det(M\psi)|^{-1} \tag{3.18}$$

The assertion of the proposition now follows from Eq. (3.14), Eq. (3.18) and the following equation:

$$\frac{(2\pi)^{d_2}}{Z} = \frac{(2\pi)^{d/2}}{Z} \overset{(*)}{=} \det{}^{\frac{1}{2}}(iS_\mu) \overset{(**)}{=} \det(iS_\mu)^{1/2} \overset{(***)}{=} \det((M\psi)^t M\psi)^{1/2} = |\det(M\psi)| \tag{3.19}$$

Here step $(*)$ follows from Example 3.4 and the assumption that $d\mu$ is normalized, step $(**)$ follows because[30] for each eigenvalue λ of S_μ also $-\lambda$ is an eigenvalue of S_μ and has the same multiplicity as λ, and step $(***)$ follows because after making the identification $V_2 \cong_\psi V_1$ we have $S_\mu = -\begin{pmatrix} 0 & M\psi \\ (M\psi)^t & 0 \end{pmatrix}$.

---

[29] a formal proof of Eq. (3.18) can be obtained after a suitable change of variable and the application of the dominated convergence theorem, cf. the proof of Eq. (3.23) below which generalizes Eq. (3.18)

[30] this can be seen, e.g., from the equation $J^{-1}S_\mu J = -S_\mu$ where $J := \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}$, 1 denoting both the identity in $\mathrm{End}(V_1)$ and $\mathrm{End}(V_2)$

□

Convention 2 *For a continuous function $f : V_0 \to \mathbb{C}$ on a d_0-dimensional Euclidean space V_0 we set*

$$\int_{V_0}^{\sim} f(x_0)dx_0 := \tfrac{1}{\pi^{d_0/2}} \lim_{\epsilon \to 0} \epsilon^{d_0/2} \int_{V_0} e^{-\epsilon|x_0|^2} f(x_0)dx_0 \tag{3.20}$$

provided that the expression on the RHS of the previous equation is well-defined. Here dx_0 is the normalized Lebesgue measure on V_0.

Proposition 3.12 *Assume that $V = V_0 \oplus V_1 \oplus V_2$ where V_0, V_1, V_2 are pairwise orthogonal subspaces of V. For each $j = 0, 1, 2$ we denote the V_j-component of $x \in V$ by x_j. Moreover, let $d\mu$ be a (centered) normalized oscillatory Gauss-type measure on $(V, \langle\cdot,\cdot\rangle)$ of the form $d\mu(x) = \frac{1}{Z}\exp(i\langle x_2, Mx_1\rangle dx$ for some linear isomorphism $M : V_1 \to V_2$. Then for every fixed $v \in V_2$ and every bounded uniformly continuous function $F : V_0 \oplus V_1 \to \mathbb{C}$ the LHS of the following equation exists iff the RHS exists and in this case we have*

$$\int_{\sim} F(x_0 + x_1)\exp(i\langle x_2, v\rangle)d\mu(x) = \int_{V_0}^{\sim} F(x_0 - M^{-1}v)dx_0 \tag{3.21}$$

where dx_0 is the normalized Lebesgue measure on V_0.

Proof. We set $d_j := \dim(V_j)$ for $j = 0, 1, 2$. Similarly as in Eq. (3.14) and with $\delta_\epsilon : V_2 \to \mathbb{R}$ as above we obtain

$$\begin{aligned} &\int_{\sim} F(x_0 + x_1)\exp(i\langle x_2, v\rangle)d\mu(x) \\ &\quad = \lim_{\epsilon\to 0}\left(\tfrac{\epsilon}{\pi}\right)^{d_0/2} \int_{V_0}\left[\int_{V_1}\left[\int_{V_2} e^{-\epsilon(|x_0|^2+|x_1|^2+|x_2|^2)} F(x_0 + x_1)\exp(i\langle x_2, v\rangle)\times\right.\right. \\ &\qquad\qquad \left.\left. \times \tfrac{1}{Z}\exp(i\langle x_2, Mx_1\rangle dx_2\right] dx_1\right] dx_0 \\ &\quad = \tfrac{1}{\pi^{d_0/2}} \lim_{\epsilon\to 0} \epsilon^{d_0/2} \int_{V_0} dx_0 e^{-\epsilon|x_0|^2}\left[\tfrac{(2\pi)^{d_2}}{Z}\int_{V_1} dx_1 e^{-\epsilon|x_1|^2} F(x_0 + x_1)\delta_\epsilon(v + Mx_1)\right]. \end{aligned} \tag{3.22}$$

Let $\psi : V_2 \to V_1$ be a fixed isometry. From the assumption that $d\mu$ was normalized it follows – using the same argument as in Eq. (3.19) above – that $\frac{(2\pi)^{d_2}}{Z} = \det^{\frac{1}{2}}(i(S_\mu)_{|V_1\oplus V_2}) = |\det(M\psi)|$. Eq. (3.22) above and the equality just mentioned will therefore imply the assertion of the proposition provided that we can show

$$\lim_{\epsilon\to 0} T(\epsilon) = 0 \tag{3.23}$$

where

$$T(\epsilon) := \epsilon^{d_0/2}\int_{V_0} dx_0 e^{-\epsilon|x_0|^2}\left[F(x_0 - M^{-1}v) - |\det(M\psi)|\int_{V_1} dx_1 e^{-\epsilon|x_1|^2} F(x_0 + x_1)\delta_\epsilon(v + Mx_1)\right] \tag{3.24}$$

In order to prove (3.23) recall that $\psi_*(dx_2) = dx_1$ and that $1 = \frac{1}{\pi^{d_2/2}} \int_{V_2} e^{-|y_2|^2} dy_2$ so we obtain

$$\begin{aligned} T(\epsilon) &= \epsilon^{d_0/2} \int_{V_0} dx_0 e^{-\epsilon|x_0|^2} \Bigg[\tfrac{1}{\pi^{d_2/2}} \int_{V_2} dy_2 e^{-|y_2|^2} F(x_0 - M^{-1}v) \\ &\qquad - \ |\det(M\psi)| \int_{V_2} dx_2 e^{-\epsilon|\psi x_2|^2} F(x_0 + \psi x_2) \tfrac{1}{(4\epsilon\pi)^{d_2/2}} e^{-\frac{|v+M\psi x_2|^2}{4\epsilon}} \Bigg] \\ &\overset{(*)}{=} \frac{1}{\pi^{d_2/2}} \int_{V_0 \oplus V_2} dy_0 dy_2 e^{-|y_0|^2 - |y_2|^2} \Big[F(\tfrac{y_0}{\sqrt{\epsilon}} - M^{-1}v) - \gamma_\epsilon(y_2) F(\tfrac{y_0}{\sqrt{\epsilon}} - M^{-1}v + \sqrt{\epsilon} 2 M^{-1} y_2) \Big] \end{aligned} \tag{3.25}$$

In Step $(*)$ we have made the changes of variable $\sqrt{\epsilon} x_0 \to y_0$ and $\frac{1}{\sqrt{\epsilon}} \frac{1}{2}(v + M\psi x_2) \to y_2$ and we have set $\gamma_\epsilon(y_2) := e^{-\epsilon|M^{-1}(2\sqrt{\epsilon} y_2 - v)|^2}$. Relation (3.23) now follows by applying the dominated convergence theorem to the last expression in Eq. (3.25) and taking into account that for all fixed $y_0 \in V_0$ and $y_2 \in V_2$ we have

$$\begin{aligned} &\lim_{\epsilon \to 0} \big[F(\tfrac{y_0}{\sqrt{\epsilon}} - M^{-1}v) - \gamma_\epsilon(y_2) F(\tfrac{y_0}{\sqrt{\epsilon}} - M^{-1}v + \sqrt{\epsilon} 2 M^{-1} y_2) \big] \\ &\quad = \lim_{\epsilon \to 0} \big[F(\tfrac{y_0}{\sqrt{\epsilon}} - M^{-1}v)(1 - \gamma_\epsilon(y_2)) \big] \\ &\qquad\qquad + \lim_{\epsilon \to 0} \gamma_\epsilon(y_2) \big[F(\tfrac{y_0}{\sqrt{\epsilon}} - M^{-1}v) - F(\tfrac{y_0}{\sqrt{\epsilon}} - M^{-1}v + \sqrt{\epsilon} 2 M^{-1} y_2) \big] = 0 + 0 = 0 \end{aligned}$$

since $\lim_{\epsilon \to 0} \gamma_\epsilon(y_2) = 1$ and, by assumption, F is bounded and uniformly continuous.

□

The following remark will be useful in Sec. 5.4.3 and Sec. 5.4.4 in [28].

Remark 3.13 If Γ is a lattice in V and $f : V \to \mathbb{C}$ a Γ-periodic continuous function then $\int_V^\sim f(x)dx$ exists and we have

$$\int_V^\sim f(x)dx = \frac{1}{vol(Q)} \int_Q f(x)dx \tag{3.26}$$

with $Q := \{\sum_i x_i e_i \mid 0 \le x_i \le 1 \forall i \le d\}$ where $(e_i)_{i \le d}$ is any fixed basis of the lattice Γ and where $vol(Q)$ denotes the volume of Q. Observe that Eq. (3.26) implies

$$\forall y \in V : \quad \int_V^\sim f(x)dx = \int_V^\sim f(x+y)dx \tag{3.27}$$

4 A "simplicial" differential geometric framework

As a preparation for Sec. 5 below we will now describe (a modified version of) some of the constructions of the "simplicial" framework in [1, 2] which will be relevant for us.

4.1 Chains and cochains

Let $d \in \mathbb{N}$ and let $\mathcal{K}$ be an oriented d-dimensional simplicial complex. For $p \in \{0, 1, \ldots, d\}$ we will denote by $\mathfrak{F}_p(\mathcal{K})$ the set of p-faces in $\mathcal{K}$. For every real finite-dimensional vector space V and every $p \in \{0, 1, \ldots, d\}$ we will denote by $C_p(\mathcal{K}, V)$ the space of "p-chains of $\mathcal{K}$ with coefficients in V" and by $C^p(\mathcal{K}, V)$ the space of "p-cochains with values in V"[31]. As usual we will denote by $\partial_{\mathcal{K}} : C_p(\mathcal{K}, V) \to C_{p-1}(\mathcal{K}, V)$, $p \in \{1, 2, \ldots, d\}$, the corresponding "boundary operator" and by $d_{\mathcal{K}} : C^p(\mathcal{K}, V) \to C^{p+1}(\mathcal{K}, V)$, $p \in \{0, 2, \ldots, d-1\}$ the corresponding "coboundary operator".

---

[31] recall that $C^p(\mathcal{K}, V)$ is simply the space of all maps $\mathfrak{F}_p(\mathcal{K}) \to V$

If $\mathcal{K}$ is finite then we can make the identification $C_p(\mathcal{K}, V) \cong C^p(\mathcal{K}, V)$ for each $p \in \{0, 1, 2, \ldots, d\}$. Moreover, if we equip V with a scalar product $\langle \cdot, \cdot \rangle$ then this induces a scalar product $\langle \cdot, \cdot \rangle_p$ on $C_p(\mathcal{K}, V) \cong C^p(\mathcal{K}, V)$ in the obvious way. Observe that the linear maps $d_{\mathcal{K}} : C^p(\mathcal{K}, V) \to C^{p+1}(\mathcal{K}, V)$ and $\partial_{\mathcal{K}} : C_{p+1}(\mathcal{K}, V) \to C_p(\mathcal{K}, V)$ are dual to each other w.r.t. the scalar products $\langle \cdot, \cdot \rangle_p$ and $\langle \cdot, \cdot \rangle_{p+1}$ i.e. we have

$$\langle d_{\mathcal{K}} \alpha, \beta \rangle_{p+1} = \langle \alpha, \partial_{\mathcal{K}} \beta \rangle_p \tag{4.1}$$

for all $\alpha \in C_p(\mathcal{K}, V) \cong C^p(\mathcal{K}, V)$ and $\beta \in C_{p+1}(\mathcal{K}, V) \cong C^{p+1}(\mathcal{K}, V)$.

Convention 3 *i) For each $p \in \{0, 1, \ldots, d\}$ we identify $\mathfrak{F}_p(\mathcal{K})$ with its image in $C^p(\mathcal{K}, \mathbb{R}) \cong C_p(\mathcal{K}, \mathbb{R})$ under the injection $\mathfrak{F}_p(\mathcal{K}) \ni \alpha \mapsto \delta_\alpha \in C^p(\mathcal{K}, \mathbb{R})$ where $\delta_\alpha \in C^p(\mathcal{K}, \mathbb{R})$ is given by $\delta_\alpha(\alpha') = \delta_{\alpha, \alpha'}$ for all $\alpha' \in \mathfrak{F}_p(\mathcal{K})$*

ii) Instead of $C_p(\mathcal{K}, \mathbb{R})$ we will write simply $C_p(\mathcal{K})$ in the following.

The constructions and definitions above can be generalized in a straightforward way to the situation where instead of a (finite) oriented simplicial complex $\mathcal{K}$ we work with a (finite) oriented "polyhedral cell complex" $\mathcal{P}$, i.e. a cell complex which is obtained by glueing together bounded "convex polytopes" in an analogous way as simplicial complexes arise from glueing together simplices, cf. part C of the Appendix for the formal definitions.

Polyhedral cell complexes arise naturally, e.g. as the duals and products of simplicial complexes (cf. Remark C.10 in part C of the Appendix).

4.2 Discrete Hodge star operators

Recall that above we fixed an oriented compact surface Σ. For the rest of this paper we will now fix two finite polyhedral cell decompositions $\mathcal{C}$ and $\mathcal{C}'$ of Σ, which are dual to each other[32], cf. Definition C.9 and the examples in Fig. 5 in part C of the Appendix and in Fig. 1 below.

Example 4.1 *In the situation in Fig. 1 below, $\mathcal{C}$ (resp. $\mathcal{C}'$) is a "hexagonal" (resp. "triangular") (polyhedral) cell decomposition. The 2-face F is dual to the 0-face x', the 0-face x is dual to the 2-face F', and the two 1-faces e and e' are dual to each other.*

Thus, in the notation of Definition C.9 in the Appendix we could write $x' = \check{F}$ or $F = \check{x}'$ and so on.

Let $\mathcal{K}$ and $\mathcal{K}'$ denote the polyhedral cell complexes corresponding to $\mathcal{C}$ and $\mathcal{C}'$ (i.e. $\mathcal{K} = (\Sigma, \mathcal{C})$ and $\mathcal{K}' = (\Sigma, \mathcal{C}')$) and let us equip $\mathcal{K}$ and $\mathcal{K}'$ with the orientations which are induced by the orientation on Σ. We can then define, for every fixed real finite-dimensional vector space V and $p \in \{0, 1, 2\}$, the operator $\star_{\mathcal{K}} : C^p(\mathcal{K}, V) \to C^{2-p}(\mathcal{K}', V)$ as the unique linear isomorphism such that for every $\alpha \in \mathfrak{F}_p(\mathcal{K}) \subset C_p(\mathcal{K})$ and $v \in V$ we have[33]

$$\star_{\mathcal{K}}(\alpha \otimes v) = \begin{cases} \check{\alpha} \otimes v & \text{if } \alpha \times \check{\alpha} \subset \mathbb{R}^p \times \mathbb{R}^{2-p} \cong \mathbb{R}^2 \text{ is "positively oriented"} \\ -\check{\alpha} \otimes v & \text{otherwise} \end{cases}$$

where $\check{\alpha} \in \mathfrak{F}_{2-p}(\mathcal{K}')$ is the face dual to α and where we have made the identifications $C^p(\mathcal{K}, V) \cong C_p(\mathcal{K}) \otimes V$ and $C^{2-p}(\mathcal{K}', V) \cong C_{2-p}(\mathcal{K}') \otimes V$.

(The operator $\star_{\mathcal{K}'} : C^p(\mathcal{K}', V) \to C^{2-p}(\mathcal{K}, V)$ will be defined in a completely analogous way.). Above we have equipped $\alpha \times \check{\alpha} \subset \mathbb{R}^2$ with the product of the orientations on α and $\check{\alpha}$. If this orientation coincides with the orientation inherited from the standard orientation of $\mathbb{R}^2$, we say that $\alpha \times \check{\alpha}$ is "positively oriented".

---

[32] for example, we can choose $\mathcal{C}$ to a be finite triangulation of Σ and $\mathcal{C}'$ to be the canonical dual of $\mathcal{C}$, cf. Remark C.10 in part C of the Appendix.

[33] recall that according to Convention 3 above by $v \cdot \alpha$ we actually mean $v \cdot \delta_\alpha$



Figure 1:

It is not difficult to see (cf. [1]) that we have

$$\partial_{\mathcal{K}'} \cdot \star_{\mathcal{K}} = \star_{\mathcal{K}} \cdot d_{\mathcal{K}} \tag{4.2}$$

and

$$\star_{\mathcal{K}}^{-1} = (-1)^{p(2-p)} \star_{\mathcal{K}'} = \begin{cases} -\star_{\mathcal{K}'} & \text{if } p = 1 \\ \star_{\mathcal{K}'} & \text{if } p \in \{0, 2\} \end{cases} \tag{4.3}$$

Moreover, when V is equipped with a scalar product and the spaces $C^p(\mathcal{K}, V)$, $C^{2-p}(\mathcal{K}', V)$, $C^p(\mathcal{K}', V)$, and $C^{2-p}(\mathcal{K}, V)$ with the induced scalar products then $\star_{\mathcal{K}}$ and $\star_{\mathcal{K}'}$ will be isometries.

Convention 4 *In the special case where $V = \mathbb{R}$ we will simply write $\star$ instead of $\star_{\mathcal{K}}$.*

(Observe that $\star\alpha$, for $\alpha \in \mathfrak{F}_p(\mathcal{K}) \subset C_p(\mathcal{K}) \cong C^p(\mathcal{K})$, then just coincides, up to a sign, with the dual face $\check{\alpha} \in \mathfrak{F}_{n-p}(\mathcal{K}') \subset C_{n-p}(\mathcal{K}) \cong C^{n-p}(\mathcal{K}')$).

Example 4.2 *In the situation in Fig. 1 we have for one*[34] *of the two possible choices of the orientation on $\mathbb{R}^2$*

$$\star x = F', \quad \star x' = F, \quad \star e = e', \quad \star e' = -e, \quad \star F = x', \quad \star F' = x,$$

cf. Convention 3 and Convention 4 above.

4.3 Discrete curves, loops, and links

Let $\mathcal{P}$ be a fixed oriented polyhedral cell complex[35]. We will call the elements of $\mathfrak{F}_0(\mathcal{P})$ (resp. $\mathfrak{F}_1(\mathcal{P})$ resp. $\mathfrak{F}_1(\mathcal{P}) \cup \{0\} \cup (-\mathfrak{F}_1(\mathcal{P})) \subset C_1(\mathcal{P})$) the "vertices" (resp. "edges" resp. "generalized edges") in $\mathcal{P}$. The element $0 \in C_1(\mathcal{P})$ will be called "the empty edge". The starting point (resp. endpoint) of an edge $e \in \mathfrak{F}_1(\mathcal{P})$ will be denoted by $\text{start}(e)$ (resp. $\text{end}(e)$), or simply by $\bullet\, e$ (resp. $e\, \bullet$).

---

[34] for the other choice of orientation the third and fourth equation have to be replaced by $\star e = -e'$ and $\star e' = e$

[35] the only five cases which will be relevant later are $\mathcal{P} \in \{\mathcal{K}, \mathcal{K}', \mathbb{Z}_N, \mathcal{K} \times \mathbb{Z}_N, \mathcal{K}' \times \mathbb{Z}_N\}$ with $\mathbb{Z}_N$ as at the beginning of Sec. 4.4

A "discrete curve" in $\mathcal{P}$ is a finite sequence $x = (x^{(k)})_{k \leq n}$, $n \in \mathbb{N}$, of vertices in $\mathcal{P}$ such that for every $k \leq n$ the two vertices $x^{(k)}$ and $x^{(k+1)}$ either coincide or are the two endpoints of an edge $e^{(k)} \in \mathfrak{F}_1(\mathcal{P})$. If $x^{(n)} = x^{(1)}$ we will call $x = (x^{(k)})_{k \leq n}$ a "discrete loop" in $\mathcal{P}$.

Every discrete curve $x = (x^{(k)})_{k \leq n}$ with $n > 1$ induces a sequence $(e^{(k)})_{k \leq n-1}$ of generalized edges given by

$$e^{(k)} = \begin{cases} e & \text{if } x^{(k)} = \text{start}(e) \\ 0 & \text{if } x^{(k)} = x^{(k+1)} \\ -e & \text{if } x^{(k)} = \text{end}(e) \end{cases}$$

where $e \in \mathfrak{F}_1(\mathcal{P})$ is the unique edge with $\{\text{start}(e), \text{end}(e)\} = \{x^{(k)}, x^{(k+1)}\}$. In the following we will mostly[36] take the "generalized edge point of view", i.e. when we mention discrete curves we consider them as sequences $e = (e^{(k)})_{k \leq n}$, $n \in \mathbb{N}$, of generalized edges and write $\bullet\, e^{(k)}$ for the corresponding vertex $x^{(k)}$. More precisely, we have[37]

$$\bullet\, e^{(k)} = \begin{cases} \text{start}(e^{(k)}) & \text{if } e^{(k)} \in \mathfrak{F}_1(\mathcal{P}), \\ \text{end}(-e^{(k)}) & \text{if } e^{(k)} \in -\mathfrak{F}_1(\mathcal{P}) \\ \text{end}(e^{(j)}) & \text{if } e^{(k)} = 0 \text{ (with } j := \max\{i \leq n \mid e^{(i)} \neq 0\}) \end{cases}$$

A discrete loop in $\mathcal{P}$ is a discrete curve $l = (l^{(k)})_{k \leq n}$ in $\mathcal{P}$ such that $\bullet\, l^{(1)} = l^{(n)}\, \bullet$

Observe that every discrete loop $l = (l^{(k)})_{k \leq n}$ in a polyhedral cell complex $\mathcal{P} = (X, \mathcal{C})$ induces, in an obvious way, a continuous loop $[0,1] \to X$, which will also be denoted by l.

A discrete link in $\mathcal{P}$ is a finite tuple $L = (l_1, l_2, \ldots, l_m)$, $m \in \mathbb{N}$, of discrete loops in $\mathcal{P}$ such that the corresponding tuple of continuous loops in X is a link in X[38].

4.4 Framings

For the rest of this paper let us fix a natural number N and we will denote by $\mathbb{Z}_N$ the cyclic group of order N. We will identify $\mathbb{Z}_N$ with the obvious subgroup of the Lie group S^1. Clearly, the points in $\mathbb{Z}_N$ induce a (polyhedral) cell decomposition of S^1 and the corresponding (1-dimensional polyhedral) cell complex will be denoted also by $\mathbb{Z}_N$ in the following[39]. We will equip S^1 with the standard (= positive) orientation. Clearly, $\mathbb{Z}_N$ can then be considered as an oriented graph.

Let $l = (l^{(k)})_{k \leq n}$ be a discrete loop in $\mathcal{K} \times \mathbb{Z}_N$ and let $l_\Sigma = (l^{(k)}_\Sigma)_{k \leq n}$ and $l_{S^1} = (l^{(k)}_{S^1})_{k \leq n}$ denote the obvious[40] "projected" discrete loops in $\mathcal{K}$ and $\mathbb{Z}_N$. According to the convention at the end of Sec. 4.3 the corresponding continuous loops in $\Sigma \times S^1$, Σ, S^1, respectively, will also be denoted by l, l_Σ, l_{S^1}.

Definition 4.3 A discrete loop $l' = (l'^{(k)})_{k \leq n'}$ in the polyhedral cell complex $\mathcal{K}' \times \mathbb{Z}_N$ is called a "framing of $l = (l^{(k)})_{k \leq n}$" iff the corresponding continuous loops l and l' in $\Sigma \times S^1$ are homotopic to each other and do not intersect.

---

[36] note that these two points of view are almost equivalent the exception being the case where $x = (x^{(k)})_{k \leq n}$ is constant. In this (degenerate) case $x = (x^{(k)})_{k \leq n}$ can of course not be recovered from $e = (e^{(k)})_{k \leq n}$ since an empty edge has now endpoints. So in this special case we are forced to take the vertex point of view

[37] observe that if $e^{(k)}$ is a "proper" edge, the notation $\bullet\, e^{(k)}$ is consistent with the notation introduced above

[38] in particular, each l_i neither intersects itself nor any of the other l_j

[39] in words: we have equipped $\mathbb{Z}_N$ with a graph structure; the set of edges is given by $\{(t, t+1) \mid t \in \mathbb{Z}_N\}$

[40] "obvious" at least in the "vertex point of view" of a discrete curve, cf. Sec. 4.3

Remark 4.4 The notion of "framing" used in [1] is slightly different. We could have made our terminology closer to the one in [1] by defining a framing of a discrete loop in $\mathcal{P} = \mathcal{K} \times \mathbb{Z}_N$ as a discrete loop (with certain additional properties) in the canonical dual $\mathcal{P}'$ of $\mathcal{P}$, cf. Remark C.10 in part C of the Appendix. However, the difference between this modified definition and Definition 4.3 above is not essential since we can identify $\mathcal{P}'$ with $\mathcal{K}' \times \mathbb{Z}_N$ in a straightforward way. More precisely: we can make the identification $\mathcal{P}' = (\mathcal{K} \times \mathbb{Z}_N)' \cong \mathcal{K}' \times \mathbb{Z}'_N \cong \mathcal{K}' \times \mathbb{Z}_N$ where in the last step we used the graph isomorphism between $\mathbb{Z}_N$ and $\mathbb{Z}'_N$ given by $\mathfrak{F}_0(\mathbb{Z}_N) \ni t \mapsto \bullet e(t) \in \mathfrak{F}_0(\mathbb{Z}'_N)$ where $e(t) = \star t \in \mathfrak{F}_1(\mathbb{Z}'_N)$.

5 Rigorous realization of $\mathrm{WLO}(L)$ by discretization: Approach I

The two approaches ("Approach I" and "Approach II") for making rigorous sense of Eq. (2.53) which we will introduce in the present section and in [28] were partly inspired by Adams' approach in [1, 2] for discretizing Abelian Chern-Simons theory or, more precisely, Abelian BF_3-theory. In fact, a crucial step in [1, 2] was the transition to the "BF-theory point of view", which involves, among other things, a suitable "field doubling", cf. Sec. 7 and Appendix C in [28].

Adams's results seem to suggest that for the discretization of *Non-Abelian* CS theory a similar strategy will have to be used. It turns out, however, that for non-Abelian CS theory the advantages of the "BF_3-theory point of view" are not as obvious as in the Abelian case (cf. Remark 7.2 in Sec. 7 of [28]) and that for the derivation of our main result, i.e. Theorem 7.3 in [28], it will be sufficient to work with the original "CS theory point of view". Accordingly, we will postpone the transition to BF_3-theory to Sec. 7 in [28].

Let us now describe the first of the two approaches mentioned above.

5.1 Definition of the spaces $\mathcal{B}(K)$, $\mathcal{A}_\Sigma(K)$ and $\mathcal{A}^\perp(K)$

Let us first introduce suitable finite-dimensional analogues of the spaces $\mathcal{B}$, $\mathcal{A}_\Sigma$ and $\mathcal{A}^\perp$.

Recall that above we fixed two finite polyhedral cell decompositions $\mathcal{C}$ and $\mathcal{C}'$ of Σ, which are assumed to be dual to each other, and we denoted by $\mathcal{K}$ and $\mathcal{K}'$ the corresponding polyhedral cell complexes. Moreover, we fixed a finite triangulation of (the Lie group) S^1 having the finite cyclic group $\mathbb{Z}_N$, $N \in \mathbb{N}$, with the obvious graph structure as the corresponding simplicial complex. Instead of $\mathcal{K}$ (resp. $\mathcal{K}'$) we will often write K_1 (resp. K_2) in the following and we set $K := (K_1, K_2)$.

For the transition from the continuum to the discrete we now make the following replacements (cf. Remark 5.1 below for an alternative definition of $\mathcal{A}^\perp(K)$):

$$\begin{array}{lcl}
S^1 & \longrightarrow & \mathbb{Z}_N \\
\Sigma & \longrightarrow & K_1 \text{ or } K_2 \\
\mathcal{B} = C^\infty(\Sigma, \mathfrak{t}) = \Omega^0(\Sigma, \mathfrak{t}) & \longrightarrow & \mathcal{B}(K) := C^0(K_1, \mathfrak{t}) \oplus C^0(K_2, \mathfrak{t}) \\
\mathcal{A}_\Sigma = \Omega^1(\Sigma, \mathfrak{g}) & \longrightarrow & \mathcal{A}_\Sigma(K) := C^1(K_1, \mathfrak{g}) \oplus C^1(K_2, \mathfrak{g}) \\
\mathcal{A}^\perp \cong C^\infty(S^1, \mathcal{A}_\Sigma) & \longrightarrow & \mathcal{A}^\perp(K) := \mathrm{Map}(\mathbb{Z}_N, \mathcal{A}_\Sigma(K))
\end{array}$$

Clearly, the scalar product $\langle \cdot, \cdot \rangle_{\mathfrak{g}}$ on $\mathfrak{g}$ induces scalar products $\ll \cdot, \cdot \gg_{\mathcal{B}(K)}$ and $\ll \cdot, \cdot \gg_{\mathcal{A}_\Sigma(K)}$ on $\mathcal{B}(K)$ and $\mathcal{A}_\Sigma(K)$ in the obvious way. We introduce a scalar product $\ll \cdot, \cdot \gg_{\mathcal{A}^\perp(K)}$ on $\mathcal{A}^\perp(K) = \mathrm{Map}(\mathbb{Z}_N, \mathcal{A}_\Sigma(K))$ by[41]

$$\ll A^\perp_1, A^\perp_2 \gg_{\mathcal{A}^\perp(K)} = \tfrac{1}{2N} \sum\nolimits_{t \in \mathbb{Z}_N} \ll A^\perp_1(t), A^\perp_2(t) \gg_{\mathcal{A}_\Sigma(K)} \tag{5.1}$$

for all $A^\perp_1, A^\perp_2 \in \mathcal{A}^\perp(K)$. In the following we will make the identification

$$\mathcal{A}^\perp(K) \cong \mathrm{Map}(\mathbb{Z}_N, C^1(K_1, \mathfrak{g})) \oplus \mathrm{Map}(\mathbb{Z}_N, C^1(K_2, \mathfrak{g})) \tag{5.2}$$

---

[41] cf. Remark 5.1 below for the factor $\frac{1}{2N}$

and we will denote by $B_j \in C^0(K_j, \mathfrak{t})$, $(A_c)_j \in C^1(K_j, \mathfrak{g})$, and $A^\perp_j \in \operatorname{Map}(\mathbb{Z}_N, C^1(K_j, \mathfrak{g}))$ (for $j \in \{1,2\}$) the obvious components of $B \in \mathcal{B}(K)$, $A_c \in \mathcal{A}_\Sigma(K)$, and $A^\perp \in \mathcal{A}^\perp(K)$.

Remark 5.1 Instead of the ansatz $\mathcal{A}^\perp(K) := \operatorname{Map}(\mathbb{Z}_N, \mathcal{A}_\Sigma(K))$ above we could use the ansatz $\mathcal{A}^\perp(K) := \mathcal{A}^\perp_{altern}(K)$ where $\mathcal{A}^\perp_{altern}(K)$ is as in Appendix D below. The latter ansatz would make it clear that we really are working in a "single field setting" rather than a "double field setting" like the one appearing in the BF_3-theory point of view (cf. the beginning of Sec. 5 above and Sec. 7 in [28]). Moreover, the use of $\mathcal{A}^\perp_{altern}(K)$ would also have several other advantages, for example, it would naturally explain the appearance of the factor $\frac{1}{2N}$ on the RHS of Eq. (5.1) above, cf. Appendix D.

In spite of these advantages we decided to work with the ansatz $\mathcal{A}^\perp(K) := \operatorname{Map}(\mathbb{Z}_N, \mathcal{A}_\Sigma(K))$ in the present paper because this simplifies the notation.

Convention 5 *For $B \in \mathcal{B}(K)$ and $x \in \mathfrak{F}_0(K_1) \cup \mathfrak{F}_0(K_2)$ and $A_c \in \mathcal{A}_\Sigma(K)$ and $e \in \mathfrak{F}_1(K_1) \cup \mathfrak{F}_1(K_2)$ we set*

$$B(x) := B_j(x) \quad \text{where } j \in \{1,2\} \text{ is given by } x \in \mathfrak{F}_0(K_j) \tag{5.3}$$

$$A_c(e) := (A_c)_j(e) \quad \text{where } j \in \{1,2\} \text{ is given by } e \in \mathfrak{F}_1(K_j) \tag{5.4}$$

Convention 6 *Let us introduce the notation*

$$\mathfrak{F}_0(K_1|K_2) := \{\bar{e} \mid e \in \mathfrak{F}_1(K_1)\} = \{\bar{e} \mid e \in \mathfrak{F}_1(K_2)\} \subset \Sigma \tag{5.5}$$

where $\bar{e}$ is the unique crossing point[42] of e and $\star e$, cf. Definition C.9 iii) in part C of the Appendix.

For $\bar{e} \in \mathfrak{F}_0(K_1|K_2)$ and $j = 1,2$ we will denote the unique element $e_j \in \mathfrak{F}_1(K_j)$ with $\bar{e} = \bar{e_j}$ by $e_j(\bar{e})$.

Definition 5.2 *Let V be a finite-dimensional real vector space.*

i) For each $x \in \mathbb{Z}_N$ we will denote by τ_x the translation operator $\operatorname{Map}(\mathbb{Z}_N, V) \to \operatorname{Map}(\mathbb{Z}_N, V)$ given by $(\tau_x f)(t) = f(t+x)$ for all $t \in \mathbb{Z}_N$.

ii) We define the operators $\hat{\partial}^{(N)}_t$, $\check{\partial}^{(N)}_t$, and $\bar{\partial}^{(N)}_t$ on $\operatorname{Map}(\mathbb{Z}_N, V)$ by

$$\hat{\partial}^{(N)}_t := N(\tau_1 - \tau_0), \tag{5.6a}$$

$$\check{\partial}^{(N)}_t := N(\tau_0 - \tau_{-1}), \tag{5.6b}$$

$$\bar{\partial}^{(N)}_t := \tfrac{N}{2}(\tau_1 - \tau_{-1}) \tag{5.6c}$$

(where 1 and -1 are the obvious elements of $\mathbb{Z}_N = \mathbb{Z}/N\mathbb{Z}$).

Clearly[43], the three difference operators $\hat{\partial}^{(N)}_t$, $\check{\partial}^{(N)}_t$ and $\bar{\partial}^{(N)}_t$ are discrete analogues of the differential operator $\partial_t := \frac{\partial}{\partial t}$ on $C^\infty(S^1, V)$.

---

[42] so in the special case where K_2 is the canonical dual of K_1 then $\bar{e}$ will be the barycenter of e

[43] this becomes totally evident by making the identification $\mathbb{Z}_N \cong \{0, \frac{1}{N}, \frac{2}{N}, \ldots, \frac{N-1}{N}\}$. Then we have $(\hat{\partial}^{(N)}_t A^\perp)(t) = \frac{A^\perp(t+\frac{1}{N}) - A^\perp(t)}{\frac{1}{N}}$, $(\check{\partial}^{(N)}_t A^\perp)(t) = \frac{A^\perp(t) - A^\perp(t-\frac{1}{N})}{\frac{1}{N}}$, and $(\bar{\partial}^{(N)}_t A^\perp)(t) = \frac{A^\perp(t+\frac{1}{N}) - A^\perp(t-\frac{1}{N})}{\frac{2}{N}}$ for $t \in \{\frac{1}{N}, \frac{2}{N}, \ldots, \frac{N-2}{N}\}$

5.2 Definition of $S_{CS}^{disc}(A^{\perp}, B)$

Let us now introduce a discrete analogue for

$$S_{CS}(A^{\perp}, B) = \pi k \big[\ll A^{\perp}, \star\big(\tfrac{\partial}{\partial t} + \operatorname{ad}(B)\big) A^{\perp} \gg_{\mathcal{A}^{\perp}} - 2 \ll A^{\perp}, \star dB \gg_{\mathcal{A}^{\perp}} \big], \tag{5.7}$$

cf. Eq. (2.47) above. In order to do so we have to look for suitable discrete analogues of the mappings $\star : \mathcal{A}_{\Sigma} \to \mathcal{A}_{\Sigma}$, $\star : C^{\infty}(S^1, \mathcal{A}_{\Sigma}) \to C^{\infty}(S^1, \mathcal{A}_{\Sigma})$, $d : C^{\infty}(\Sigma, \mathfrak{g}) \to \mathcal{A}_{\Sigma}$, the scalar product $\ll \cdot, \cdot \gg_{\mathcal{A}^{\perp}}$ on $\mathcal{A}^{\perp} \cong C^{\infty}(S^1, \mathcal{A}_{\Sigma})$ appearing in Eq. (5.7), and the differential operator $\frac{\partial}{\partial t}$.

For the transition from the continuum setting to the discrete setting let us now make the following replacements (using the identification (5.2)):

$$\begin{array}{lcl}
\ll \cdot, \cdot \gg_{\mathcal{A}^{\perp}} & \longrightarrow & \ll \cdot, \cdot \gg_{\mathcal{A}^{\perp}(K)} \\[2ex]
\star & \longrightarrow & \begin{pmatrix} 0 & \star_{K_2} \\ \star_{K_1} & 0 \end{pmatrix} =: \star_K \\
d & \longrightarrow & \begin{pmatrix} d_{K_1} & 0 \\ 0 & d_{K_2} \end{pmatrix} =: d_K \\
\frac{\partial}{\partial t} & \longrightarrow & \begin{pmatrix} \hat{\partial}_t^{(N)} & 0 \\ 0 & \check{\partial}_t^{(N)} \end{pmatrix} =: \partial_t^{(N)}
\end{array}$$

Here $d_{K_j} : C^0(K_j, \mathfrak{t}) \to C^1(K_j, \mathfrak{t})$, $j = 1, 2$, are as in Sec. 4. The operators $\star_{K_1} : \operatorname{Map}(\mathbb{Z}_N, C^1(K_1, \mathfrak{g})) \to \operatorname{Map}(\mathbb{Z}_N, C^1(K_2, \mathfrak{g}))$, $\star_{K_2} : \operatorname{Map}(\mathbb{Z}_N, C^1(K_2, \mathfrak{g})) \to \operatorname{Map}(\mathbb{Z}_N, C^1(K_1, \mathfrak{g}))$ are the linear isomorphisms defined in the obvious way[44]. Finally, the difference operator $\hat{\partial}_t^{(N)}$ (resp. $\check{\partial}_t^{(N)}$) is given by Definition 5.2 above with $V = C^1(K_1, \mathfrak{g})$ (resp. $V = C^1(K_2, \mathfrak{g})$).

Remark 5.3 Regarding the definition of the operator $\partial_t^{(N)}$ above one might wonder why instead of the operators $\bar{\partial}_t^{(N)}$ and $\check{\partial}_t^{(N)}$ we did not use the (anti-)symmetrized version(s) $\bar{\partial}_t^{(N)} : \operatorname{Map}(\mathbb{Z}_N, C^1(K_j, \mathfrak{g})) \to \operatorname{Map}(\mathbb{Z}_N, C^1(K_j, \mathfrak{g}))$, $j = 1, 2$, cf. Definition 5.2 above. This point will be clarified in part D of the Appendix below.

Using the replacements listed above we now arrive at the following discrete analogue for $S_{CS}(A^{\perp}, B)$:

$$S_{CS}^{disc}(A^{\perp}, B) := \pi k \Big[\ll A^{\perp}, \star_K \big(\partial_t^{(N)} + \operatorname{ad}(B)\big) A^{\perp} \gg_{\mathcal{A}^{\perp}(K)} - 2 \ll A^{\perp}, \star_K d_K B \gg_{\mathcal{A}^{\perp}(K)} \Big] \tag{5.8}$$

for $A^{\perp} \in \mathcal{A}^{\perp}(K)$, $B \in \mathcal{B}(K)$. Here $\operatorname{ad}(B) : \mathcal{A}^{\perp}(K) \to \mathcal{A}^{\perp}(K)$ is given by[45]

$$(\operatorname{ad}(B) A^{\perp})(t)(e) = \operatorname{ad}(B(\bar{e})) \cdot A^{\perp}(t)(e) \tag{5.9}$$

for all $A^{\perp}$, $t \in \mathbb{Z}_N$, and $e \in \mathfrak{F}_1(K_1) \cup \mathfrak{F}_1(K_2)$ where we have set for $\bar{e} \in \mathfrak{F}_0(K_1|K_2)$

$$B(\bar{e}) := \tfrac{1}{2}[B_1(\bar{e}) + B_2(\bar{e})] \tag{5.10}$$

Here we have extended for each $j \in \{1, 2\}$ the field $B_j : \mathfrak{F}_0(K_j) \to \mathfrak{t}$ to a map on the bigger set $\mathfrak{F}_0(K_j) \cup \mathfrak{F}_0(K_1|K_2)$ by setting

$$B_j(\bar{e_j}) := \tfrac{1}{2}[B_j(\bullet\, e_j) + B_j(e_j\, \bullet)] \qquad \forall e_j \in \mathfrak{F}_1(K_j) \tag{5.11}$$

Observation 5.4 *The operators* $\star_K \partial_t^{(N)} : \mathcal{A}^{\perp}(K) \to \mathcal{A}^{\perp}(K)$ *and* $\star_K \operatorname{ad}(B) : \mathcal{A}^{\perp}(K) \to \mathcal{A}^{\perp}(K)$ *are symmetric w.r.t to* $\ll \cdot, \cdot \gg_{\mathcal{A}^{\perp}(K)}$.

---

[44] i.e. by $(\star_{K_j} A_j^{\perp})(t) = \star_{K_j}(A_j^{\perp}(t))$ for each $A_j^{\perp} \in \operatorname{Map}(\mathbb{Z}_N, C^1(K_j, \mathfrak{g}))$, $t \in \mathbb{Z}_N$, and $j = 1, 2$ where $\star_{K_j} : C^1(K_j, \mathfrak{g}) \to C^1(K_{3-j}, \mathfrak{g})$ is given as in Sec. 4 above

[45] Note that if we had made the ansatz $(\operatorname{ad}(B) A^{\perp})(t)(e) = \operatorname{ad}(B(e)) \cdot A^{\perp}(t)(e)$ with $B(e) := \frac{1}{2}\big(B(\bullet\, e) + B(e\, \bullet)\big)$ instead of the ansatz in Eq. (5.9) above the operator $\operatorname{ad}(B) : \mathcal{A}^{\perp}(K) \to \mathcal{A}^{\perp}(K)$ would in general not be antisymmetric w.r.t. $\ll \cdot, \cdot \gg_{\mathcal{A}^{\perp}(K)}$ (and, consequently, the operator $\star_K \operatorname{ad}(B)$ would not be symmetric). In fact, the operator $\operatorname{ad}(B)$ given by Eq. (5.9) is exactly the operator which is obtained by anti-symmetrizing the operator $\operatorname{ad}(B)$ appearing at the beginning of the present footnote

5.3 Definition of $\mathrm{Hol}^{disc}_l(A^\perp, B; \mathrm{h})$

Let us introduce a discrete analogue $\mathrm{Hol}^{disc}_l(A^\perp, B; \mathrm{h})$ for the expressions $\mathrm{Hol}_l(A^\perp, B; \mathrm{h}) = \mathrm{Hol}_l(A^\perp + A_{\mathrm{sg}}(\mathrm{h}) + Bdt)$ appearing in Eq. (2.54c) above.

5.3.1 Unframed version

We start with the observation that for $A^\perp \in \mathcal{A}^\perp \cong C^\infty(S^1, \mathcal{A}_\Sigma)$ and $B \in \mathcal{B} = C^\infty(\Sigma, \mathfrak{t})$ we have

$$A^\perp(l'(t)) = A^\perp(l_{S^1}(t))(l'_\Sigma(t)), \qquad (Bdt)(l'(t)) = B(l_\Sigma(t)) \cdot dt(l'_{S^1}(t)) \tag{5.12}$$

for $t \in [0,1]$. Here we have set $l_\Sigma := \pi_\Sigma \circ l$ and $l_{S^1} := \pi_{S^1} \circ l$ where $\pi_\Sigma : \Sigma \times S^1 \to \Sigma$, $\pi_{S^1} : \Sigma \times S^1 \to S^1$ are the canonical projections. From Eqs. (2.4), (2.54c), and (5.12) we obtain

$$\begin{aligned} &\mathrm{Hol}_l(A^\perp, B; \mathrm{h}) \\ &= \lim_{n\to\infty} \prod_{k=1}^n \exp\Big(\tfrac{1}{n}\big(A^\perp(l_{S^1}(t))(l'_\Sigma(t)) + A_{\mathrm{sg}}(\mathrm{h})(l'_\Sigma(t)) + B(l_\Sigma(t)) \cdot dt(l'_{S^1}(t))\big)\Big)_{|t=\frac{k}{n}} \end{aligned} \tag{5.13}$$

Let us now discretize the RHS of this equation. Let $l = (l^{(k)})_{k\le n}$ be a discrete loop in $K_1 \times \mathbb{Z}_N$ and let $(l^{(k)}_\Sigma)_{k\le n}$ and $(l^{(k)}_{S^1})_{k\le n}$ denote the projected discrete loops in K_1 and $\mathbb{Z}_N$, cf. the paragraph before Definition 4.3 above.

In contrast to the situation in the continuum setting where the parameter $n \in \mathbb{N}$ was sent to ∞ we will leave n fixed. It is now natural to make the replacements

$$\begin{array}{lll} l_\Sigma(\frac{k}{n}) & \longrightarrow & \bullet\, l^{(k)}_\Sigma \\ l_{S^1}(\frac{k}{n}) & \longrightarrow & \bullet\, l^{(k)}_{S^1} \\ \frac{1}{n} l'_\Sigma(\frac{k}{n}) & \longrightarrow & l^{(k)}_\Sigma \\ \frac{1}{n} l'_{S^1}(\frac{k}{n}) & \longrightarrow & l^{(k)}_{S^1} \\ dt & \longrightarrow & dt^{(N)} \end{array}$$

with $dt^{(N)} \in C^1(\mathbb{Z}_N, \mathbb{R})$ given by $dt^{(N)}(e) = \frac{1}{N}$ for all $e \in \mathfrak{F}_1(\mathbb{Z}_N)$. We will make the identification $C^1(\mathbb{Z}_N, \mathbb{R}) \cong \mathrm{Hom}_{\mathbb{R}}(C_1(\mathbb{Z}_N), \mathbb{R})$, so for a generalized edge $e \in \mathfrak{F}_1(\mathbb{Z}_N) \cup \{0\} \cup (-\mathfrak{F}_1(\mathbb{Z}_N)) \subset C_1(\mathbb{Z}_N)$ we then have $dt^{(N)}(e) = \frac{1}{N}\,\mathrm{sgn}(e)$ where $\mathrm{sgn}(e) \in \{-1, 0, 1\}$ is given by $\mathrm{sgn}(e) := \pm 1$ if $e \in \pm\mathfrak{F}_1(\mathbb{Z}_N)$ and $\mathrm{sgn}(e) := 0$ if $e = 0$.

Applying the replacements above to the RHS of Eq. (5.13) we arrive at the ansatz

$$\mathrm{Hol}^{disc}_l(A^\perp, B; \mathrm{h}) := \prod_{k=1}^n \exp\Big(A^\perp(\bullet\, l^{(k)}_{S^1})(l^{(k)}_\Sigma) + A_{\mathrm{sg}}(\mathrm{h})(l^{(k)}_\Sigma) + B(\bullet\, l^{(k)}_\Sigma) \cdot \tfrac{1}{N}\,\mathrm{sgn}(l^{(k)}_{S^1})\Big) \tag{5.14}$$

for $A^\perp \in \mathcal{A}^\perp(K)$, $B \in \mathcal{B}(K)$, $\mathrm{h} \in [\Sigma, G/T]$ where

$$A_{\mathrm{sg}}(\mathrm{h})(l^{(k)}_\Sigma) := \int_{l^{(k)}_\Sigma} A_{\mathrm{sg}}(\mathrm{h}) \tag{5.15}$$

where on the right-hand side of Eq. (5.15) we have identified $l^{(k)}_\Sigma$ with the obvious smooth curve $c : [0,1] \to \Sigma$ induced by $l^{(k)}_\Sigma$.

5.3.2 Framed version

Instead of working with the simple discrete loop l in $K_1 \times \mathbb{Z}_N$ we can work with a suitable pair (l, l') where l' is a "framing" of l in the sense of Sec. 4.4. Without loss of generality[46] we can assume that l and l' have the same length n.

---

[46] since we can always add "empty" edges to l or l' if necessary

We can then introduce the following framed analogue of Eq. (5.14) above

$$\begin{aligned}\mathrm{Hol}_l^{disc}(A^{\perp}, B; \mathrm{h}) := \prod\nolimits_{k=1}^{n} \exp\Big(&\tfrac{1}{2}\big(A^{\perp}(\bullet\, l_{S^1}^{(k)})\big)(l_{\Sigma}^{(k)}) + \tfrac{1}{2}\big(A^{\perp}(\bullet\, l_{S^1}^{'(k)})\big)(l_{\Sigma}^{'(k)}) \\ &+ \tfrac{1}{2}\big(A_{\mathrm{sg}}(\mathrm{h})(l_{\Sigma}^{(k)}) + A_{\mathrm{sg}}(\mathrm{h})(l_{\Sigma}^{'(k)})\big) + \tfrac{1}{2}B(\bullet\, l_{\Sigma}^{(k)}) \cdot \tfrac{1}{N}\,\mathrm{sgn}(l_{S^1}^{(k)}) + \tfrac{1}{2}B(\bullet\, l_{\Sigma}^{'(k)}) \cdot \tfrac{1}{N}\,\mathrm{sgn}(l_{S^1}^{'(k)})\Big) \end{aligned} \tag{5.16}$$

Remark 5.5 As a preparation for [28] let us mention that if we set $\mathrm{Hol}_l^{disc}(\check{A}^{\perp}, A_c, B; \mathrm{h}) := \mathrm{Hol}_l^{disc}(\check{A}^{\perp} + A_c, B; \mathrm{h})$ for fixed $\check{A}^{\perp} \in \check{\mathcal{A}}^{\perp}(K)$, $A_c \in \mathcal{A}_c^{\perp}(K)$, $B \in \mathcal{B}(K)$, and $\mathrm{h} \in [\Sigma, G/T]$ (where $\check{\mathcal{A}}^{\perp}(K)$ and $\mathcal{A}_c^{\perp}(K)$ are as in Sec. 5.7 below) then we have

$$\begin{aligned}\mathrm{Hol}_l^{disc}(\check{A}^{\perp}, A_c, B; \mathrm{h}) = \prod\nolimits_{k} \exp\Big(&\tfrac{1}{2}\big(\check{A}^{\perp}(\bullet\, l_{S^1}^{(k)})\big)(l_{\Sigma}^{(k)}) + \tfrac{1}{2}\big(\check{A}^{\perp}(\bullet\, l_{S^1}^{'(k)})\big)(l_{\Sigma}^{'(k)}) + \tfrac{1}{2}A_c(l_{\Sigma}^{(k)}) + \tfrac{1}{2}A_c(l_{\Sigma}^{'(k)}) \\ &+ \frac{1}{2}\int_{l_{\Sigma}^{(k)}} A_{\mathrm{sg}}(\mathrm{h}) + \frac{1}{2}\int_{l_{\Sigma}^{'(k)}} A_{\mathrm{sg}}(\mathrm{h}) + \tfrac{1}{2}B(\bullet\, l_{\Sigma}^{(k)}) \cdot \tfrac{1}{N}\,\mathrm{sgn}(l_{S^1}^{(k)}) + \tfrac{1}{2}B(\bullet\, l_{\Sigma}^{'(k)}) \cdot \tfrac{1}{N}\,\mathrm{sgn}(l_{S^1}^{'(k)})\Big) \end{aligned} \tag{5.17}$$

The last formula will be useful in [28].

5.4 Definition of $\mathrm{Det}_{FP}^{disc}(B)$

We will need a discrete analogue $\mathrm{Det}_{FP}^{disc}(B)$ of the factor $\mathrm{Det}_{FP}(B) = \det\big(1_{\mathfrak{k}} - \exp(\mathrm{ad}(B))_{|\mathfrak{k}}\big)$ appearing in Eq. (2.54) above. We make the ansatz

$$\mathrm{Det}_{FP}^{disc}(B) := \prod\nolimits_{x \in \mathfrak{F}_0(K_1) \cup \mathfrak{F}_0(K_2)} \det\big(1_{\mathfrak{k}} - \exp(\mathrm{ad}(B(x)))_{|\mathfrak{k}}\big) \tag{5.18}$$

for every $B \in \mathcal{B}(K)$.

5.5 Discrete version of $\int_{\Sigma \backslash \{\sigma_0\}} \mathrm{Tr}\big(dA_{\mathrm{sg}}(\mathrm{h}) \cdot B\big)$

Next we introduce a suitable discrete analogue of the expression $\int_{\Sigma \backslash \{\sigma_0\}} \mathrm{Tr}\big(dA_{\mathrm{sg}}(\mathrm{h}) \cdot B\big)$, namely the expression

$$\int_{\Sigma \backslash \{\sigma_0\}} \mathrm{Tr}\big(dA_{\mathrm{sg}}(\mathrm{h}) \cdot \overline{B}\big) \tag{5.19}$$

where we have set for every $B = (B_1, B_2) \in \mathcal{B}(K) = C^0(K_1, \mathfrak{t}) \oplus C^0(K_2, \mathfrak{t})$

$$\overline{B} := \frac{1}{2}[\chi_1(B_1) + \chi_2(B_2)] \in \mathrm{Map}(\Sigma, \mathfrak{t}) \tag{5.20}$$

Here $\chi_j : C^0(K_j, \mathfrak{t}) \to \mathrm{Map}(\Sigma, \mathfrak{t})$, $j \in \{1, 2\}$, are the two mappings given by

$$\chi_j(B_j) := \sum\nolimits_{F \in \mathfrak{F}_2(K_j)} B_j(\check{F}) \cdot 1_F \tag{5.21}$$

with 1_F denoting the indicator function of the subset $F \subset \Sigma$ and $\check{F} = \star F$.

We will assume in the following that σ_0 does not lie in the 1-skeleton of K_1 or of K_2.

5.6 Discrete version of $1_{C^\infty(\Sigma,\mathfrak{t}_{reg})}(B)$

Let us introduce a discrete analogue of the indicator function $1_{C^\infty(\Sigma,\mathfrak{t}_{reg})}(B)$. The obvious candidate is $\prod_{x \in \mathfrak{F}_0(K_1)\cup\mathfrak{F}_0(K_2)} 1_{\mathfrak{t}_{reg}}(B(x))$. However, with this choice we would get some problems later due to the fact that $1_{\mathfrak{t}_{reg}} : \mathfrak{t} \to \{0,1\} \subset [0,1]$ is non-continuous. For this reason we will regularize the function $1_{\mathfrak{t}_{reg}}$. We fix a family $(1^{(s)}_{\mathfrak{t}_{reg}})_{s>0}$, of elements of $C^\infty_{\mathbb{R}}(\mathfrak{t})$, with the following properties:

- $\text{Image}(1^{(s)}_{\mathfrak{t}_{reg}}) \subset [0,1] \quad \text{and} \quad \text{supp}(1^{(s)}_{\mathfrak{t}_{reg}}) \subset \mathfrak{t}_{reg}$
- $1^{(s)}_{\mathfrak{t}_{reg}} \to 1_{\mathfrak{t}_{reg}}$ pointwise as $s \to 0$
- Each $1^{(s)}_{\mathfrak{t}_{reg}}$ is invariant under the operation of the affine Weyl group $\mathcal{W}_{\text{aff}}$ on $\mathfrak{t}$, cf. part A of the Appendix below.

For fixed $s > 0$ we take as the discrete analogue of $1_{C^\infty(\Sigma,\mathfrak{t}_{reg})}(B)$ the expression

$$\prod_x 1^{(s)}_{\mathfrak{t}_{reg}}(B(x)) := \prod_{x \in \mathfrak{F}_0(K_1)\cup\mathfrak{F}_0(K_2)} 1^{(s)}_{\mathfrak{t}_{reg}}(B(x)) \tag{5.22}$$

Later we will let $s \to 0$.

5.7 Discrete version of the decomposition $\mathcal{A}^\perp = \check{\mathcal{A}}^\perp \oplus \mathcal{A}^\perp_c$

We introduce a decomposition of $\mathcal{A}^\perp(K)$, which is analogous to the decomposition $\mathcal{A}^\perp = \check{\mathcal{A}}^\perp \oplus \mathcal{A}^\perp_c$ in Sec. 2.3.3. In order to do so we introduce the notation

$$\mathcal{A}_{\Sigma,V}(K) := C^1(K_1, V) \oplus C^1(K_2, V) \tag{5.23}$$

for every real vector space V. We then have the following analogue of the decomposition $\mathcal{A}^\perp = \check{\mathcal{A}}^\perp \oplus \mathcal{A}^\perp_c$ in Sec. 2.3.3 above namely,

$$\mathcal{A}^\perp(K) = \check{\mathcal{A}}^\perp(K) \oplus \mathcal{A}^\perp_c(K) \tag{5.24}$$

where

$$\mathcal{A}^\perp_c(K) := \{A^\perp \in \mathcal{A}^\perp(K) \mid A^\perp \text{ is constant and } \mathcal{A}_{\Sigma,\mathfrak{t}}(K)\text{-valued}\} \cong \mathcal{A}_{\Sigma,\mathfrak{t}}(K) \tag{5.25a}$$

$$\check{\mathcal{A}}^\perp(K) := \{A^\perp \in \mathcal{A}^\perp(K) \mid \sum_{t \in \mathbb{Z}_N} A^\perp(t) \in \mathcal{A}_{\Sigma,\mathfrak{k}}(K)\} \tag{5.25b}$$

Observe that $\mathcal{A}_\Sigma(K) = \mathcal{A}_{\Sigma,\mathfrak{g}}(K) \cong \mathcal{A}_{\Sigma,\mathfrak{t}}(K) \oplus \mathcal{A}_{\Sigma,\mathfrak{k}}(K)$ and that for every $\check{A}^\perp \in \check{\mathcal{A}}^\perp(K)$, $A^\perp_c \in \mathcal{A}^\perp_c(K)$, and $B \in \mathcal{B}(K)$ we have

$$S^{disc}_{CS}(\check{A}^\perp + A^\perp_c, B) = S^{disc}_{CS}(\check{A}^\perp, B) + S^{disc}_{CS}(A^\perp_c, B) \tag{5.26}$$

with the two expressions on the RHS given explicitly by Eqs. (5.27) and (5.29) below.

For referring to the elements of the space $\check{\mathcal{A}}^\perp(K)$ we will use the variable $\check{A}^\perp$ and for the elements of the space $\mathcal{A}_{\Sigma,\mathfrak{t}}(K) \cong \mathcal{A}^\perp_c(K)$ the variable $A^\perp_c$ (and later sometimes the variable A_c).

5.8 Discrete versions of the two Gauss-type measures in Eq. (2.53)

Clearly, we have

$$S^{disc}_{CS}(\check{A}^\perp, B) = \pi k \ll \check{A}^\perp, \star_K \big(\partial^{(N)}_t + \text{ad}(B)\big) \cdot \check{A}^\perp \gg_{\mathcal{A}^\perp(K)} \tag{5.27}$$

for all $\check{A}^{\perp} \in \check{\mathcal{A}}^{\perp}(K)$ and $B \in \mathcal{B}(K)$. Moreover, if $B \in \mathcal{B}(K)$ fulfills $\prod_x 1_{\mathfrak{t}_{reg}}(B(x)) \neq 0$ then $\partial_t^{(N)} + \mathrm{ad}(B) : \check{\mathcal{A}}^{\perp}(K) \to \check{\mathcal{A}}^{\perp}(K)$ will be injective provided that N is sufficiently large. In this case the (rigorous) complex measure

$$\exp(iS^{disc}_{CS}(\check{A}^{\perp}, B))D\check{A}^{\perp} \tag{5.28}$$

is a non-degenerate centered oscillatory Gauss type measure on $\check{\mathcal{A}}^{\perp}(K)$. Here we have equipped $\check{\mathcal{A}}^{\perp}(K)$ with the restriction $\ll \cdot, \cdot \gg_{\check{\mathcal{A}}^{\perp}(K)}$ of the scalar product $\ll \cdot, \cdot \gg_{\mathcal{A}^{\perp}(K)}$ onto $\check{\mathcal{A}}^{\perp}(K)$ and $D\check{A}^{\perp}$ denotes the normalized Lebesgue measure on $\check{\mathcal{A}}^{\perp}(K)$ w.r.t. $\ll \cdot, \cdot \gg_{\check{\mathcal{A}}^{\perp}(K)}$.

Moreover, since

$$S^{disc}_{CS}(A^{\perp}_c, B) = -2\pi k \ll A^{\perp}_c, \star_K d_K B \gg_{\mathcal{A}^{\perp}(K)} \tag{5.29}$$

the complex measure

$$\nu := \exp(iS^{disc}_{CS}(A^{\perp}_c, B))(DA^{\perp}_c \otimes DB) \tag{5.30}$$

is a (degenerate) centered oscillatory Gauss type measure on $\mathcal{A}^{\perp}_c(K) \times \mathcal{B}(K)$. Here we have equipped $\mathcal{A}^{\perp}_c(K) \times \mathcal{B}(K)$ with the scalar product $\ll \cdot, \cdot \gg_{\mathcal{A}^{\perp}_c(K) \oplus \mathcal{B}(K)} := \ll \cdot, \cdot \gg_{\mathcal{A}^{\perp}_c(K)} \oplus \ll \cdot, \cdot \gg_{\mathcal{B}(K)}$ and $DA^{\perp}_c$ and DB denote the obvious normalized Lebesgue measures.

5.9 Definition of $\mathrm{WLO}^{disc}_{rig}(L)$

Let $\mathcal{K}$ and $N \in \mathbb{N}$ be as above. For the rest of this paper we will fix a discrete link $L = (l_1, l_2, \ldots, l_m)$ in $\mathcal{K} \times \mathbb{Z}_N = K_1 \times \mathbb{Z}_N$ with "colors" $(\rho_1, \rho_2, \ldots, \rho_m)$, $m \in \mathbb{N}$. Moreover, for each l_i we will fix a "framing" l'_i, i.e. a suitable discrete loop in $\mathcal{K}' \times \mathbb{Z}_N = K_2 \times \mathbb{Z}_N$ fulfilling the conditions in Sec. 4.4. Using the definitions of the previous subsections[47] we then arrive at the following simplicial analogue $\mathrm{WLO}^{disc}_{rig}(L)$ of the heuristic expression $\mathrm{WLO}(L)$ in Eq. (2.53)

$$\begin{aligned}
\mathrm{WLO}^{disc}_{rig}(L) := \lim_{s \to 0} \sum\nolimits_{\mathrm{h} \in [\Sigma, G/T]} \int_{\sim} \big(\prod\nolimits_x 1^{(s)}_{\mathfrak{t}_{reg}}(B(x))\big) \, \mathrm{Det}^{disc}_{FP}(B) \\
\times \Big[\int_{\sim} \prod\nolimits_{i=1}^{m} \mathrm{Tr}_{\rho_i}\big(\mathrm{Hol}^{disc}_{l^i}(\check{A}^{\perp} + A^{\perp}_c, B; \mathrm{h})\big) \exp(iS^{disc}_{CS}(\check{A}^{\perp}, B)) D\check{A}^{\perp} \Big] \\
\times \exp\big(i\tfrac{k}{2\pi} \int_{\Sigma \backslash \{\sigma_0\}} \mathrm{Tr}\big(dA_{\mathrm{sg}}(\mathrm{h}) \cdot \overline{B}\big)\big) \exp(iS^{disc}_{CS}(A^{\perp}_c, B))(DA^{\perp}_c \otimes DB)
\end{aligned} \tag{5.31}$$

5.10 Three modifications

Naively, one might expect that – at least for the simple type of links and framings mentioned in Sec. 2.3.3 above – the expression $\mathrm{WLO}^{disc}_{rig}(L)$ is well-defined and that we have

$$\mathrm{WLO}^{disc}_{rig}(L) \sim |L| \tag{5.32}$$

where $|L|$ is the shadow invariant (associated to G and k) of the link L, cf. (2.55) above and Appendix in B in [28].

It turns out, however, that we can only get a result in this direction if we modify our original approach. In order to do so we will now make three modifications (Mod1)–(Mod3). Additionally, we will either have to work with a new discretization $L^{(N)}(B)$ of the continuum operator $\frac{\partial}{\partial t} + \mathrm{ad}(B)$ appearing in Eq. (5.7) above or we will have to perform a continuum limit "in the S^1-component", cf. Remark 5.9 below.

Remark 5.6 i) Modification (Mod2) will be eliminated[48] in Approach II in [28] and Modification (Mod1) will be replaced by a more natural condition in Approach II.

---

[47] by $\mathrm{Hol}^{disc}_{l^i}(\check{A}^{\perp} + A^{\perp}_c, B; \mathrm{h}))$ in Eq. (5.31) we mean the "framed version" as given by Eq. (5.16) in Subsec. 5.3.2

[48] cf., however, Sec. 3.4 in [28]

ii) Modification (Mod3) is rather artificial and should not be taken too seriously. It can be eliminated by performing the transition to the BF-theory point of view mentioned in Sec. 5 above. In order to avoid long formulas, which would make our computations even heavier, we will postpone the transition to the BF-theory point of view to the end of [28]. The main part of [28] deals with Approach II which is based on the original "single field point of view" and which also allows the elimination of (Mod3).

Modification (Mod1)

Let us now reconsider the question of what a suitable discrete analogue $\mathrm{Det}^{disc}_{FP}(B)$ of the continuum expression $\mathrm{Det}_{FP}(B) = \det\big(1_{\mathfrak{k}} - \exp(\mathrm{ad}(B))_{|\mathfrak{k}}\big)$ should be. Above we made the ansatz

$$\mathrm{Det}^{disc}_{FP}(B) = \prod\nolimits_{x \in \mathfrak{F}_0(K_1) \cup \mathfrak{F}_0(K_2)} \det\big(1_{\mathfrak{k}} - \exp(\mathrm{ad}(B(x)))_{|\mathfrak{k}}\big) \tag{5.33}$$

We will now modify Eq. (5.33) and make the ansatz

$$\begin{aligned}\mathrm{Det}^{disc}_{FP}(B) := \prod\nolimits_{x \in \mathfrak{F}_0(K_1) \cup \mathfrak{F}_0(K_2)} & \det\big(1_{\mathfrak{k}} - \exp(\mathrm{ad}(B(x)))_{|\mathfrak{k}}\big)^{1/2} \\ & \times \prod\nolimits_{\bar{e} \in \mathfrak{F}_0(K_1|K_2)} \big(\det\big(1_{\mathfrak{k}} - \exp(\mathrm{ad}(B(\bar{e})))_{|\mathfrak{k}}\big)^{1/2}\end{aligned} \tag{5.34}$$

instead.

Remark 5.7 The expression $\det\big(1_{\mathfrak{k}} - \exp(\mathrm{ad}(b))_{|\mathfrak{k}}\big)$ is a "natural square" i.e. it is of the from $T(b)^2$ where $T(b)$ is "nice". Thus the two square roots appearing on the RHS of Eq. (5.34) just cancel the 2-exponents.

Modification (Mod2)

We define $B(\bar{e})$ for $\bar{e} \in \mathfrak{F}_0(K_1|K_2)$ not by $B(\bar{e}) := \frac{1}{2}[B_1(\bar{e}) + B_2(\bar{e})]$ as in Eq. (5.10) above but by

$$B(\bar{e}) := \begin{cases} B_1(\bar{e}) & \text{if } B_1(\bullet\, e_1) = B_1(e_1 \bullet) \text{ and } B_2(\bullet\, e_2) \neq B_2(e_2 \bullet) \\ B_2(\bar{e}) & \text{if } B_2(\bullet\, e_2) = B_2(e_2 \bullet) \text{ and } B_1(\bullet\, e_1) \neq B_1(e_1 \bullet) \\ \frac{1}{2}[B_1(\bar{e}) + B_2(\bar{e})] & \text{otherwise} \end{cases} \tag{5.35}$$

where $e_1 := e_1(\bar{e})$ and $e_2 := e_2(\bar{e})$, cf. Convention 6 in Sec. 5.1.

Modification (Mod3)

In Eq. (5.31) we replace[49] the space $\mathcal{B}(K) = B_c(K) \oplus \mathcal{B}'(K)$ where

$$\begin{aligned}\mathcal{B}_c(K) &:= \{B \in \mathcal{B}(K) \mid \ B_1 \text{ and } B_2 \text{ are constant}\} \cong \mathfrak{t}^2 \\ \mathcal{B}'(K) &:= \{B \in \mathcal{B}(K) \mid B_1(\sigma_1) = B_2(\sigma_2) = 0\}\end{aligned}$$

by the slightly smaller subspace $B_{cc}(K) \oplus \mathcal{B}'(K)$ where

$$B_{cc}(K) := \{B \in \mathcal{B}_c(K) \mid B_1 = B_2\} \cong \mathfrak{t} \tag{5.36}$$

Above $\sigma_1 \in \mathfrak{F}_0(K_1)$ and $\sigma_2 \in \mathfrak{F}_0(K_2)$ are[50] any two points which do not lie on $\mathrm{arc}(l^i_\Sigma) \cup \mathrm{arc}(l'^i_\Sigma)$, $i \leq m$, and which are "close" to each other in the sense that there is some $F \in \mathfrak{F}_2(K_1) \cup \mathfrak{F}_2(K_2)$ with $\sigma_1, \sigma_2 \in \overline{F}$.

---

[49] In particular, the complex measure $\exp(iS^{disc}_{CS}(A^\perp_c, B))(DA^\perp_c \otimes DB))$ appearing in Eq. (5.31) will no longer be the measure ν on $\mathcal{A}^\perp_c(K) \times \mathcal{B}(K)$, introduced in Eq. (5.30) but the "restriction" of ν onto the subspace $\mathcal{A}^\perp_c(K) \times (B_{cc}(K) \oplus \mathcal{B}'(K))$

[50] more generally, we could choose any two points $\sigma_1 \in \mathfrak{F}_0(K_1)$ and $\sigma_2 \in \mathfrak{F}_0(K_2)$ which lie in the same connected component of $\Sigma \backslash \bigcup_{i=1}^m \big(\mathrm{arc}(l^i_\Sigma) \cup \mathrm{arc}(l'^i_\Sigma)\big)$

5.11 Definition of $\mathrm{WLO}_{rig}(L)$

As mentioned above, the three modifications (Mod1)–(Mod3) alone are not sufficient if we want to arrive at a result of the form (5.32). That is why we will now replace the discrete operator $\partial_t^{(N)} + \mathrm{ad}(B)$ appearing in Eq. (5.8) above by another natural discretization of the continuum operator $\frac{\partial}{\partial t} + \mathrm{ad}(B)$ in Eq. (5.7), namely by the operator $L^{(N)}(B)$ which will be introduced in Sec. 3.3 of [28].

Let $\mathrm{WLO}^{mod}_{rig}(L)$ be defined as $\mathrm{WLO}^{disc}_{rig}(L)$ above but using the operator $L^{(N)}(B)$ instead of the operator $\partial_t^{(N)} + \mathrm{ad}(B)$ and using the three modifications (Mod1), (Mod2), and (Mod3) above. Setting

$$\mathrm{WLO}_{rig}(L) := \frac{\mathrm{WLO}^{mod}_{rig}(L)}{\mathrm{WLO}^{mod}_{rig}(\emptyset)} \tag{5.37}$$

where $\emptyset$ is the "empty" link[51] we then arrive at the following preliminary version of Theorem 7.3 in [28]:

Theorem 5.8 *Assume that the link $L = (l_1, l_2, \ldots, l_m)$ fixed above fulfills conditions (NCP) and (NH) of Sec. 5.1 in [28] and that the framings $(l'_1, l'_2, \ldots, l'_m)$ fulfill conditions (FC1)–(FC4) of Sec. 5.2 in [28]. Assume also that $k \geq c_{\mathfrak{g}}$ where $c_{\mathfrak{g}}$ is the dual Coxeter number of $\mathfrak{g}$. Then $\mathrm{WLO}_{rig}(L)$ is well-defined and we have*

$$\mathrm{WLO}_{rig}(L) = \frac{|L|}{|\emptyset|} \tag{5.38}$$

where $|\cdot|$ is the shadow invariant associated to G and k, cf. Remark 5.4 in [28] and Appendix B in [28] for the full definitions and for additional comments.

The derivation of Eq. (5.38) is similar to the proof of Theorem 7.3 in [28]. Firstly, the inner (oscillatory) integral $\int_\sim \cdots \exp(\cdots) D\check{A}^\perp$ in Eq. (5.31) is evaluated. For the special class of links and framings considered in the theorem above this is easy and basically boils down to an application of Proposition 3.7 (cf. Step 1 in Sec. 5.4 in [28]). One is left with an (oscillatory) integral of the form $\int_\sim \cdots \exp(\cdots)(DA_c \otimes DB)$ which only involves Abelian fields, namely B and A_c. This remaining (oscillatory) integral is evaluated using[52] Proposition 3.12 (cf. Step 2 in Sec. 5.4 in [28]). After this one proceeds in a similar way as in Steps 3–6 in Sec. 5.4 in [28].

Remark 5.9 If we want to work with the original discretization of the continuum operator $\frac{\partial}{\partial t} + \mathrm{ad}(B)$ in Eq. (5.7), i.e. with the operator $\partial_t^{(N)} + \mathrm{ad}(B)$ we need to perform a suitable continuum limit "in the S^1-component".

As a first step we set $N(n) := N \cdot n$ for every $n \in \mathbb{N}$. Observe that the link $L = (l_1, \ldots, l_m)$ in $\mathcal{K} \times \mathbb{Z}_N$ can be considered to be a link in the "finer" polyhedral cell complex $\mathcal{K} \times \mathbb{Z}_{N(n)}$ and each of the loops $l'_1, \ldots, l'_m$ can be considered to be a loop in $\mathcal{K}' \times \mathbb{Z}_{N(n)}$.

Next we rewrite Eq. (5.31) above using the identification $\mathcal{A}_c^\perp(K) \cong \mathcal{A}_{\Sigma,\mathrm{t}}(K)$ (The advantage of the space $\mathcal{A}_{\Sigma,\mathrm{t}}(K)$ is that it does not depend on $N(n)$ while the space $\mathcal{A}_c^\perp(K)$ does). Then we include a $n \to \infty$-limit "inside" the $\int_\sim \cdots (DA_c^\perp \otimes DB)$ integral (but "outside" the $\int_\sim \cdots D\check{A}^\perp$ integral) on the RHS of Eq. (5.31).

Perhaps it is also possible to perform the $n \to \infty$-limit after both integrations in Eq. (5.31) have been carried out. This would have stylistic advantages but the computations would then become unnecessarily difficult.

---

[51] so $\mathrm{WLO}^{disc}_{rig}(\emptyset)$ is a discrete analogue of the partition function $Z(\Sigma \times S^1)$

[52] or rather, the generalization of Proposition 3.12 where V_0 is not assumed to be orthogonal to V_1

Acknowledgements: I want to thank the anonymous referee of my paper [26] whose comments motivated me to look for an alternative approach for making sense of the RHS of Eq. (2.53), which is less technical than the continuum approach in [24, 26, 27]. This eventually led to the present paper and its sequel [28]. I am also grateful to Jean-Claude Zambrini for several comments which led to improvements in the presentation of the paper.

Finally, it is a great pleasure for me to thank Benjamin Himpel for many useful and important comments and suggestions, which not only had a major impact on the presentation and overall structure of the present paper and [28] but also inspired me to reconsider the issue of discretizing the operator $\partial_t + \mathrm{ad}(B)$, appearing in Eq. (5.7) above (this later led to Sec. 3.3 in [28]).

A Appendix: Lie theoretic notation I

A.1 List of notation I

- G: the simply-connected compact Lie group fixed in Sec. 2.1
- $\mathfrak{g}$: the Lie algebra of G
- T: the maximal torus of G fixed in Sec. 2.2
- $\mathfrak{t}$: the Lie algebra of T
- exp: the exponential map $\mathfrak{g} \to G$ of G.
- $G_{reg} := \{g \in G \mid g$ is regular, i.e. is contained in exactly one maximal torus of $G\}$
- $T_{reg} := T \cap G_{reg}$
- $\mathfrak{g}_{reg} := \exp^{-1}(G_{reg})$
- $\mathfrak{t}_{reg} := \exp^{-1}(T_{reg}) = \exp_{|\mathfrak{t}}^{-1}(T_{reg})$
- $\langle \cdot, \cdot \rangle$: the unique Ad-invariant scalar product on $\mathfrak{g}$ such that $\langle \check{\alpha}, \check{\alpha} \rangle = 2$ holds for every short real coroot $\check{\alpha}$ of the pair $(\mathfrak{g}, \mathfrak{t})$, cf. Appendix A in [28].
- $\mathfrak{k}$: the $\langle \cdot, \cdot \rangle$-orthogonal complement of $\mathfrak{t}$ in $\mathfrak{g}$
- $\pi_{\mathfrak{t}}$: the $\langle \cdot, \cdot \rangle$-orthogonal projection $\mathfrak{g} \to \mathfrak{t}$
- $\pi_{\mathfrak{k}}$: the $\langle \cdot, \cdot \rangle$-orthogonal projection $\mathfrak{g} \to \mathfrak{k}$
- P: the Weyl alcove fixed in Sec. 2.2.4. Recall that a Weyl alcove is simply a connected component of $\mathfrak{t}_{reg}$.
- $\mathcal{W}_{\text{aff}} \subset \mathrm{Aff}(\mathfrak{t})$: the affine Weyl group associated to $(\mathfrak{g}, \mathfrak{t})$, cf. Appendix A in [28]. We remark that $\mathcal{W}_{\text{aff}}$ operates freely and transitively on the set of Weyl alcoves.

Using $\langle \cdot, \cdot \rangle$ we can make the identification $\mathfrak{t} \cong \mathfrak{t}^*$. Sometimes we write $\langle \cdot, \cdot \rangle_{\mathfrak{g}}$ instead of $\langle \cdot, \cdot \rangle$.

A.2 Example: the special case $G = SU(2)$

Let us now consider the special group $G = SU(2)$ with the "standard" maximal torus, i.e. the

maximal torus $T = \{\exp(\theta\tau \mid \theta \in \mathbb{R}\}$ where $\tau := \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix}$

Then we have (with $\cong$ meaning "homeomorphic" and with the convention $S\tau := \{t\tau \mid t \in S\}$ for any $S \subset \mathbb{R}$):

$$\begin{aligned}
G &= SU(2) = \{A \in \mathrm{Mat}(2,\mathbb{C}) \mid AA^* = 1, \det(A) = 1\} && \cong S^3 \\
\mathfrak{g} &= su(2) = \{A \in \mathrm{Mat}(2,\mathbb{C}) \mid A + A^* = 0, \mathrm{Tr}(A) = 0\} && \cong \mathbb{R}^3 \\
T &= \{\exp(\theta\tau \mid \theta \in \mathbb{R}\} = \left\{ \begin{pmatrix} e^{i\theta} & 0 \\ 0 & e^{-i\theta} \end{pmatrix} \mid \theta \in \mathbb{R} \right\} && \cong S^1 \\
\mathfrak{t} &= \mathbb{R} \cdot \tau = \{\theta\tau \mid \theta \in \mathbb{R}\} && \cong \mathbb{R} \\
G/T &= \{gT \mid g \in G\} && \cong S^2 \\
G_{reg} &= SU(2)\backslash\{-1, 1\} && \cong S^2 \times (0,1) \\
\mathfrak{g}_{reg} &= \mathfrak{g}\backslash \bigcup\nolimits_{n \in \mathbb{N}_0} \{b \in \mathfrak{g} \mid |b| = n\} && \cong S^2 \times (\mathbb{R}_+\backslash\mathbb{N}) \\
T_{reg} &= T\backslash\{-1, 1\} && \cong S^1\backslash\{-1, 1\} \\
\mathfrak{t}_{reg} &= \mathfrak{t}\backslash\{n\pi\tau \mid n \in \mathbb{Z}\} && \cong \mathbb{R}\backslash\mathbb{Z}
\end{aligned}$$

Moreover, we have

- $\exp : \mathfrak{g} \to G$ is the restriction of the exponential map of $\mathrm{Mat}(2,\mathbb{C})$ onto $\mathfrak{g}$, i.e.

$$\exp(A) = \sum\nolimits_{n=0}^{\infty} \tfrac{A^n}{n!} \in G \subset \mathrm{Mat}(2,\mathbb{C}) \quad \text{for} \quad A \in \mathfrak{g} \subset \mathrm{Mat}(2,\mathbb{C})$$

- the scalar product $\langle \cdot, \cdot \rangle$ is given by $\langle A, B \rangle = -\frac{1}{4\pi^2} \mathrm{Tr}_{\mathrm{Mat}(2,\mathbb{C})}(A \cdot B)$ for all $A, B \in su(2)$. The norm $|\cdot|$ appearing in the formula for $\mathfrak{g}_{reg}$ denotes the norm associated to this scalar product.
- $\mathfrak{k} = \left\{ \begin{pmatrix} 0 & -z \\ \bar{z} & 0 \end{pmatrix} \mid z \in \mathbb{C} \right\}$
- the set of Weyl alcoves is $\{P_n \mid n \in \mathbb{Z}\}$ where $P_n := (n\pi, (n+1)\pi)\ \tau$. Accordingly, for P we could take, e.g., the set $P := P_0 = (0, \pi)\tau$.
- $\mathcal{W}_{\mathrm{aff}}$: the subgroup of the affine group $\mathrm{Aff}(\mathfrak{t})$ which is generated by the reflection $\mathfrak{t} \ni b \mapsto -b \in \mathfrak{t}$ and the translation $\mathfrak{t} \ni b \mapsto b + 2\pi\tau \in \mathfrak{t}$.

Let us also mention that

$$\det(1_{\mathfrak{k}} - \exp(\mathrm{ad}(x\tau))_{|\mathfrak{k}}) = 4\sin^2(x), \qquad x \in \mathbb{R}$$

B Appendix: Some technical details for Sec. 2.2.4

In Sec. 2.2.4 we omitted some technical points. We will now give some more details. In Example B.1 below we make things a bit more explicit in the special case $G = SU(2)$ (using some of the explicit relations from part A of the Appendix).

i) Let us briefly sketch why (2.23) holds. Observe first that

$$A_{reg}/\mathcal{G} \cong (A_{reg}/\tilde{\mathcal{G}})/\mathcal{G}_\Sigma \tag{B.1}$$

where $\tilde{\mathcal{G}} := \{\Omega \in \mathcal{G} \mid \sigma \in \Sigma : \Omega((\sigma, 1)) = 1\}$. Secondly, if S is a connected component of $\mathfrak{g}_{reg} = \exp^{-1}(G_{reg})$ then $\exp : S \to G_{reg}$ is a bijection[53]. It is not difficult to see that this implies that also $q : A^{\perp} \times C^{\infty}(\Sigma, S) \ni (A^{\perp}, B) \mapsto (A^{\perp} + Bdt) \cdot \tilde{\mathcal{G}} \in A_{reg}/\tilde{\mathcal{G}}$ is a bijection, cf. Proposition 3.1 in [25]. Thirdly, if P is an alcove contained in S then the map $\theta : P \times G/T \ni (b, \bar{g}) \mapsto \bar{g} b \bar{g}^{-1} \in S$ is a well-defined homeomorphism[54]. Thus we have the identification

---

[53] this follow because $\exp : \mathfrak{g}_{reg} \to G_{reg}$ is a covering and G_{reg} is simply-connected, cf. Example B.1 below for the special case $G = SU(2)$

$$\mathcal{A}_{reg}/\tilde{\mathcal{G}} \cong \mathcal{A}^{\perp} \times C^{\infty}(\Sigma, S) \cong \mathcal{A}^{\perp} \times C^{\infty}(\Sigma, P) \times C^{\infty}(\Sigma, G/T) \tag{B.2}$$

(observe that under this identification the obvious $\mathcal{G}_{\Sigma}$-operation on $\mathcal{A}_{reg}/\tilde{\mathcal{G}}$ induces the $\mathcal{G}_{\Sigma}$-operation on $\mathcal{A}^{\perp} \times C^{\infty}(\Sigma, P) \times C^{\infty}(\Sigma, G/T)$, which is given by $(A^{\perp}, B, \bar{g}) \cdot \Omega = (A^{\perp} \cdot \Omega, B, \Omega^{-1}\bar{g})$ for each $\Omega \in \mathcal{G}_{\Sigma}$).

Fourthly, observe that the map $p : \mathcal{A}^{\perp} \times C^{\infty}(\Sigma, P) \times C^{\infty}(\Sigma, G/T)/\mathcal{G}_{\Sigma} \to \big(\mathcal{A}^{\perp} \times C^{\infty}(\Sigma, P) \times C^{\infty}(\Sigma, G/T)\big)/\mathcal{G}_{\Sigma}$ which maps each $(A^{\perp}, B, \mathrm{h})$ to the $\mathcal{G}_{\Sigma}$-orbit of $(A^{\perp}, B, \bar{g}_{\mathrm{h}})$ is a surjection.

Finally, observe that under the two identifications (B.1) and (B.2) the map p coincides with the restriction of $\Pi_{\overline{V}}$ to $\mathcal{A}^{\perp} \times C^{\infty}(\Sigma, P) \times [\Sigma, G/T]$. Thus we have $\mathcal{A}_{reg}/\mathcal{G} = \Pi_{\overline{V}}(\mathcal{A}^{\perp} \times C^{\infty}(\Sigma, P) \times [\Sigma, G/T]) \subset \mathrm{Image}(\Pi_{\overline{V}})$.

ii) In the derivation in part i) we did not make use of the assumption that Σ is compact, so in fact relation (2.23) also holds for noncompact Σ. But since[55] the set $[\Sigma, G/T]$ then just consists of the single point $[1_T]$, the relation (2.23) reduces to relation (2.14) above.

iii) By contrast, if Σ is compact then relation (2.14) does *not* hold. In order to see this observe that in this case $[\Sigma, G/T]$ has infinitely many elements, cf. Remark 2.5. The map p in part i) is not injective but it does have the weaker property that $\mathrm{h}_1 \neq \mathrm{h}_2$ implies $p(A_1^{\perp}, B_1, \mathrm{h}_1) \neq p(A_2^{\perp}, B_2, \mathrm{h}_2)$. Thus for compact Σ the set

$$\Pi_{\overline{V}}(\mathcal{A}^{\perp} \times C^{\infty}(\Sigma, P) \times \{[1_T]\}) = \pi_{\mathcal{G}}(\mathcal{A}^{\perp} \oplus C^{\infty}(\Sigma, P)dt) = \pi_{\mathcal{G}}(\mathcal{A}^{qax}(T)) \cap \mathcal{A}_{reg}/\mathcal{G}$$

will be a proper subset of $\Pi_{\overline{V}}(\mathcal{A}^{\perp} \times C^{\infty}(\Sigma, P) \times [\Sigma, G/T]) = \mathcal{A}_{reg}/\mathcal{G}$, cf. part i). Clearly, this implies $\mathcal{A}_{reg}/\mathcal{G} \not\subset \pi_{\mathcal{G}}(\mathcal{A}^{qax}(T))$.

Example B.1 It is probably instructive to verify some of the claims made above (and some of the claims made in Sec. 2.2.4) directly in the special case where $\Sigma = S^2$ and where $G = SU(2)$.

i) We claimed above that G_{reg} is always simply-connected. In the special case $G = SU(2)$ this follows immediately from the concrete formula $G_{reg} \cong S^2 \times (0, 1)$ in part A of the Appendix

ii) Let S be a fixed connected component of $\mathfrak{g}_{reg}$, e.g. $S := \{b \in \mathfrak{g} \mid 0 < |b| < 1\}$, and P a Weyl alcove contained in S. Above we claimed that $\exp : S \to G_{reg}$ and $\theta : P \times G/T \ni (b, \bar{g}) \mapsto \bar{g} b \bar{g}^{-1} \in S$ are homeomorphisms. In particular, this means that $G_{reg} \cong S \cong P \times G/T$. In the special case $G = SU(2)$ the last statement follows immediately from the concrete formulas in part A of the Appendix.

iii) Recall that in the case $G = SU(2)$ the fiber bundle $\pi_{G/T} : G \to G/T$ is just the Hopf fibration. Since $\pi_2(S^3) = 0$, a topologically non-trivial (i.e. not null-homotopic) map $\bar{g} : S^2 \to S^2$ cannot be lifted to a map $\Omega : S^2 \to S^3$. On the other hand, the restriction of $\bar{g}$ to the (contractible) subset $S^2 \backslash \{\sigma_0\} \cong \mathbb{R}^2$ always admits such a lift. (This illustrates Remark 2.4 in Sec. 2.2.4 in the present special case.)

iv) Remark 2.5 in Sec. 2.2.4 implies that there is a natural bijection from $[\Sigma, G/T]$ to $I = \ker(\exp_{|\mathfrak{t}})$. In the special case $\Sigma = S^2$ we can verify[56] this directly since $[\Sigma, G/T] = [S^2, G/T] = \pi_2(G/T)$ (as sets) and $\pi_2(G/T) \cong \pi_2(S^2) \cong \mathbb{Z} \cong I$.

---

[54] cf. Example B.1 below for the special case $G = SU(2)$

[55] this follows, e.g., by combining the observation in Footnote 13 with the first two observations in Remark 2.4

[56] of course in order to check that the map in Remark 2.5 is such a bijection we will have to show more

C Appendix: Polyhedral cell complexes

C.1 Polyhedral cell decompositions and complexes

Talking informally, a "polyhedral cell-complex" is just a cell-complex which is obtained by glueing together "convex polytopes" in an analogous way as simplicial complexes arise from glueing together simplices. Here a "convex polytope" is the generalization of the notion of a convex polygon and convex polyhedron (cf. Fig. 2 for some examples) to arbitrary dimension.



Figure 2:

For the sake of completeness let us give a formal definition:

Definition C.1 *i) Let V be a finite-dimensional real vector space. A convex polytope*[57] *in V is a non-empty bounded subset P of V which is of the form*

$$P = \bigcap_{a \in A} H_a \tag{C.1}$$

where $(H_a)_{a \in A}$ is a finite family of closed halfspaces of V. Here with "closed halfspace" we mean a subset H of V of the form $H = \{x \in V \mid l(x) \geq 0\}$ for some non-trivial linear form $l : V \to \mathbb{R}$.

ii) Let V and P be as above, let $(H_a)_{a \in A}$ be a family of halfspaces such that (C.1) *holds, and let h_a be the hyperplane bounding H_a, for $a \in A$.*

A "face" of P is a non-empty proper subset S of P of the form $S = P \cap \bigcap_{a \in A'} h_a$ where $A' \subset A$. Observe that each face S of P is again a convex polytope in V (since each h_a is the intersection of the two closed half spaces which it is bounding).

iii) An abstract convex polytope (or simply, a "convex polytope") is a pair (P, V) where V is a finite-dimensional real vector space and P a convex polytope in V (equipped with the topology inherited from the standard topology of V). Instead of (P, V) we will often write simply P.

iv) The dimension $\dim(P)$ *of an abstract convex polytope $P = (P, V)$ is the dimension of the linear span*[58] *V_P of the subset $P - x$ in V where x in an arbitrary point in P.*

An orientation on $P = (P, V)$ is a non-vanishing element of $\Lambda^d V_P$ where $d := \dim(V_P)$.

---

[57] or, more precisely, a closed bounded convex polytope

[58] clearly, V_P does not depend on the special choice of x

Remark C.2 Observe that, as a topological space, an abstract convex polytope P is just a closed n-cell[59], where $n := \dim(P)$. Thus an abstract convex polytope can be considered to be a closed cell with additional structure (we will refer to this in Remark C.6 below).

The next notion which we want to introduce is the notion of a "polyhedral cell decomposition". The figures below should give a good idea of what we mean here. More precisely, Fig. 3 shows a polyhedral cell decomposition of a closed square and Fig. 4 shows a polyhedral cell decomposition of S^2. (Other examples are given in Fig. 5 below).



Figure 3:



Figure 4:

Here is a formal definition[60]:

Definition C.3 *A "polyhedral cell decomposition" $\mathcal{C}$ of a topological space X is a family $\mathcal{C} = ((P_a, \Phi_a))_{a \in A}$ where each $P_a = (V_a, P_a)$ is an (abstract) convex polytope and Φ_a is an embedding $P_a \to X$ such that the following conditions are fulfilled:*

i) Each point $x \in X$ lies in exactly one of the sets $\Phi_a(\overset{\circ}{P}_a)$, $a \in A$. Here $\overset{\circ}{P}_a$ denotes the interior of P_a in V_a.

ii) For each face F of P_a, $a \in A$, the restriction $(\Phi_a)_{|F}$ of Φ_a onto F can be "identified" with one of the Φ_b, $b \in A$. More precisely, we have $(\Phi_a)_{|F} \circ \psi_{|P_b} = \Phi_b$ for a suitably chosen $b \in A$ and a suitably chosen affine injective map $V_b \to V_a$ with the property $\psi(P_b) = F$.

iii) A set $S \subset X$ is open in X iff $\Phi_a^{-1}(S)$ is open in P_a for all a.

Remark C.4 In the terminology of Definition C.7 below, condition i) above just says that X is the disjoint union of all the "open cells" of $\mathcal{C}$. This motivates the term "cell decomposition".

Definition C.5 *A "polyhedral cell complex" is a pair $\mathcal{P} = (X, \mathcal{C})$ where X is a topological (Hausdorff) space and $\mathcal{C}$ is a "polyhedral cell decomposition" of X*

Remark C.6 i) If we recall the definition of a CW-complex we see that a polyhedral cell complex is a regular CW-complex where

1. Each of the closed cells involved has some extra structure (cf. Remark C.2 above), and,
2. The way in which these closed cells are glued together respects this extra structure (as explained in condition ii) in Definition C.3 above).

ii) Clearly, the notions "polyhedral cell decomposition" and "polyhedral cell complex" generalize the notions "triangulation" and "simplicial complex".

Definition C.7 *Let $\mathcal{C} = ((P_a, \Phi_a))_{a \in A}$ be a polyhedral cell decomposition of X and $p \in \mathbb{N}_0$.*

---

[59] i.e. homoemorphic to an n-dimensional closed ball

[60] we remark that this definition was modeled after the definition of a "$\triangle$-complex" structure in [30]

i) We set $\mathfrak{F}_p(\mathcal{C}) := \{\Phi_a(P_a) \mid a \in A$ *with* $\dim(P_a) = p\}$*. The elements of* $\mathfrak{F}_p(\mathcal{C})$ *are called the "p-faces" of* $\mathcal{C}$*.*

ii) We set $\mathrm{Cell}_p(\mathcal{C}) := \{\Phi_a(\overset{\circ}{P}_a) \mid a \in A$ *with* $\dim(P_a) = p\}$*. The elements of* $\mathrm{Cell}(\mathcal{C}) := \bigcup_p \mathrm{Cell}_p(\mathcal{C})$ *are called the "open cells" of* $\mathcal{C}$*.*

iii) If $\mathcal{P} = (X, \mathcal{C})$ *is polyhedral cell complex we write* $\mathfrak{F}_p(\mathcal{P})$ *instead of* $\mathfrak{F}_p(\mathcal{C})$*.*

Remark C.8 There is a natural 1-1-correspondence between the elements of $\mathrm{Cell}_p(\mathcal{C})$ and of $\mathfrak{F}_p(\mathcal{C})$: $E \in \mathrm{Cell}_p(\mathcal{C})$ corresponds to $F \in \mathfrak{F}_p(\mathcal{C})$ iff F is the closure of E in X

Convention 7 *Let* X *and* $\mathcal{C} = ((P_a, \Phi_a))_{a \in A}$ *be as above, let* $F \in \mathfrak{F}_p(\mathcal{K})$*,* $p \in \mathbb{N}_0$ *and let* $b \in A$ *be given by* $F = \Phi_b(P_b)$*. We then usually identify* F *with the convex polytope* P_b *via the homeomorphism* Φ_b*. Clearly, after doing so, the notions of the "barycenter"* $\bar{F}$ *of* F *and the "convex hull"* $[x, y]$ *of two points* $x, y, \in F$ *are then well-defined.*

C.2 Dual cell decompositions

Let Σ be a surface[61].

Definition C.9 *Let* $\mathcal{C}$ *and* $\mathcal{C}'$ *be two polyhedral cell decompositions of* Σ*.*

i) We say that $\mathcal{C}'$ *and* $\mathcal{C}$ *are dual to each other iff*

- *There is a bijection* $\psi_0 : \mathrm{Cell}_0(\mathcal{C}) \to \mathrm{Cell}_2(\mathcal{C}')$ *such that* $x \in \psi_0(x)$ *for all* $x \in \mathrm{Cell}_0(\mathcal{C})$*.*
- *There is a bijection* $\psi_1 : \mathrm{Cell}_1(\mathcal{C}) \to \mathrm{Cell}_1(\mathcal{C}')$ *such that each* $e \in \mathrm{Cell}_1(\mathcal{C})$ *intersects* $\psi_1(e) \in \mathrm{Cell}_1(\mathcal{C}')$ *in exactly one point and* e *intersects none of the other elements of* $\mathrm{Cell}_1(\mathcal{C}')$*.*
- *There is a bijection* $\psi_2 : \mathrm{Cell}_2(\mathcal{C}) \to \mathrm{Cell}_0(\mathcal{C}')$ *such that* $x' \in \psi_2^{-1}(x)$ *for all* $x' \in \mathrm{Cell}_0(\mathcal{C}')$*.*

Observe that each of the bijections $\psi_p : \mathrm{Cell}_p(\mathcal{C}) \to \mathrm{Cell}_{2-p}(\mathcal{C}')$ *is necessarily unique and induces a bijection* $\bar{\psi}_p : \mathfrak{F}_p(\mathcal{C}) \to \mathfrak{F}_{2-p}(\mathcal{C}')$ *(via the 1-1-correspondence in Remark C.8)*

ii) If $\mathcal{C}$ *and* $\mathcal{C}'$ *are dual to each other we set* $\check{F} := \bar{\psi}_p(F) \in \mathfrak{F}_{2-p}(\mathcal{C}')$ *for* $F \in \mathfrak{F}_p(\mathcal{C})$ *and* $\check{F}' := \bar{\psi}_p^{-1}(F') \in \mathfrak{F}_{2-p}(\mathcal{C})$ *for* $F' \in \mathfrak{F}_p(\mathcal{C}')$*.*

We call call $\check{F}$ *(resp.* $\check{F}'$*) the face "dual" to* F *(resp. "dual" to* F'*), cf. Example 4.1 in Sec. 4.2 above.*

iii) If $\mathcal{C}$ *and* $\mathcal{C}'$ *are dual to each other then for each* $e \in \mathfrak{F}_1(\mathcal{C})$ *(or* $e \in \mathfrak{F}_1(\mathcal{C}')$*) the unique intersection point of* e *and the dual face* $\check{e}$ *will be denoted by*[62] $\bar{e}$*.*

Figure 5 below, which is taken from [30], gives some examples.

Remark C.10 Finally, let us mention that if $\mathcal{C} = ((P_a, \Phi_a))_a$ is a polyhedral cell decomposition of Σ then in the set of polyhedral cell decomposition which are dual to $\mathcal{C}$ there is a distinguished element. We denote this element by $\mathcal{C}'$ and call it the canonical dual of $\mathcal{C}$, or simply, "the" dual of $\mathcal{C}$.

$\mathcal{C}'$ is constructed in a two step process where one first "breaks down" $\mathcal{C}$ into smaller pieces, which are given by the so-called "barycentric subdivision" of $\mathcal{C}$, and then, in a second step,

---

[61] We remark that Definition C.9 can be generalized to higher dimensional manifolds but we will not need this in the present paper

[62] the notation $\bar{e}$ is motivated by the fact that in the special case where $\mathcal{C}'$ is the canonical dual of $\mathcal{C}$ in the sense of Remark C.10 below, $\bar{e}$ will just be the "barycenter" of e



Figure 5:

reassembles these pieces in a suitable way. (In fact, this construction works in arbitrary dimension, and can even be used to define "the" dual $\mathcal{P}'$ of an abstract polyhedral cell complexes $\mathcal{P}'$, see, e.g. [1], for the special case where $\mathcal{P}$ is an (arbitrary) simplicial complex).

In the present case where Σ is 2-dimensional we have the following explicit[63] description of the dual $\mathcal{C}'$ (cf. Convention 7 above):

- $\mathfrak{F}_0(\mathcal{C}') := \{\bar{F} \mid F \in \mathfrak{F}_2(\mathcal{C})\}$ where $\bar{F}$ is the "barycenter" of F
- $\mathfrak{F}_1(\mathcal{C}') := \{[x_+(e), \bar{e}] \cup [\bar{e}, x_-(e)] \mid e \in \mathfrak{F}_1(\mathcal{C})\}$ where, for each $e \in \mathfrak{F}_1(\mathcal{C})$, $\bar{e}$ is the "barycenter" of e and $x_+(e)$ and $x_-(e)$ are the barycenters of the two 2-faces $F_+(e), F_-(e) \in \mathfrak{F}_2(\mathcal{C})$ which bound e.
- The elements of $\mathfrak{F}_2(\mathcal{C}')$ are the closures of the connected components of $\Sigma \backslash (\mathfrak{F}_0(\mathcal{C}') \cup \mathfrak{F}_1(\mathcal{C}'))$

D The space $\mathcal{A}^{\perp}_{altern}(K)$

Several aspects/points of Approach I (and also Approach II in [28]) get clearer[64] if, instead of working with the space

$$\mathcal{A}^{\perp}(K) \cong \mathrm{Map}(\mathbb{Z}_N, C^1(K_1, \mathfrak{g})) \oplus \mathrm{Map}(\mathbb{Z}_N, C^1(K_2, \mathfrak{g})) \tag{D.1}$$

introduced in Sec. 5.1 above, we work with the following "alternating" variant $\mathcal{A}^{\perp}_{altern}(K)$, given by

$$\mathcal{A}^{\perp}_{altern}(K) := \mathrm{Map}(\mathbb{Z}_{2N}^{odd}, C^1(K_1, \mathfrak{g})) \oplus \mathrm{Map}(\mathbb{Z}_{2N}^{even}, C^1(K_2, \mathfrak{g})) \tag{D.2}$$

where

$$\begin{aligned} \mathbb{Z}_{2N}^{even} &:= \{[t] \mid t \in \{2, 4, 6, \ldots, 2N\}\} \subset \mathbb{Z}_{2N} \\ \mathbb{Z}_{2N}^{odd} &:= \{[t] \mid t \in \{1, 3, 5, \ldots, 2N-1\}\} \subset \mathbb{Z}_{2N} \end{aligned}$$

---

[63] strictly speaking we would also have to describe the maps $\Phi'_{a'}$, $a' \in A'$, explicitly. Since this is both technical and straightforward we omit this here

[64] but the notation gets more complicated, which is why we decided to work with the space $\mathcal{A}^{\perp}(K)$ in the main text

(Here we have set $[t] := \pi(t)$ where $\pi : \mathbb{Z} \to \mathbb{Z}_{2N}$ is the canonical projection).

Observe that if we make the identifications $\mathbb{Z}_{2N}^{odd} \cong \mathbb{Z}_N \cong \mathbb{Z}_{2N}^{even}$ induced by the bijections $j_{odd} : \mathbb{Z}_N \to \mathbb{Z}_{2N}^{odd}$ and $j_{even} : \mathbb{Z}_N \to \mathbb{Z}_{2N}^{even}$ which are given by

$$j_{odd}([t]) = [2t-1], \quad j_{even}([t]) = [2t] \quad \text{for all } t \in \{1, 2, \ldots, N\} \tag{D.3}$$

we obtain an identification

$$\mathcal{A}^{\perp}_{altern}(K) \cong \mathcal{A}^{\perp}(K) \tag{D.4}$$

Thus $\mathcal{A}^{\perp}(K)$ and $\mathcal{A}^{\perp}_{altern}(K)$ are "equivalent". However, $\mathcal{A}^{\perp}_{altern}(K)$ has the following two stylistic advantages over $\mathcal{A}^{\perp}(K)$:

Advantage 1 Recall that in the definition of the scalar product $\ll \cdot, \cdot \gg_{\mathcal{A}^{\perp}(K)}$ in Eq. (5.1) a factor $\frac{1}{2N}$ appeared. This factor arises totally naturally when working with the space $\mathcal{A}^{\perp}_{altern}(K)$ instead of $\mathcal{A}^{\perp}(K)$. In this case the "natural" scalar product $\ll \cdot, \cdot \gg_{\mathcal{A}^{\perp}_{altern}(K)}$ on $\mathcal{A}^{\perp}_{altern}(K)$ will be

$$\ll A^{\perp}_1, A^{\perp}_2 \gg_{\mathcal{A}^{\perp}_{altern}(K)} = \tfrac{1}{2N} \sum\nolimits_{t \in \mathbb{Z}_{2N}} \ll A^{\perp}_1(t), A^{\perp}_2(t) \gg_{\mathcal{A}_{\Sigma}(K)}$$

$\ll \cdot, \cdot \gg_{\mathcal{A}^{\perp}(K)}$ in Eq. (5.1) is just the scalar product on $\mathcal{A}^{\perp}(K)$ obtained from $\ll \cdot, \cdot \gg_{\mathcal{A}^{\perp}_{altern}(K)}$ after making the identification (D.4).

Advantage 2 Recall that in Sec. 5.2 we defined the operator $\partial_t^{(N)} : \mathcal{A}^{\perp}(K) \to \mathcal{A}^{\perp}(K)$ by

$$\partial_t^{(N)} := \begin{pmatrix} \hat{\partial}_t^{(N)} & 0 \\ 0 & \check{\partial}_t^{(N)} \end{pmatrix} \tag{D.5}$$

where $\hat{\partial}_t^{(N)}$ and $\check{\partial}_t^{(N)}$ are as in Definition 5.2 and where the matrix notation refers to the decomposition (D.1) above.

It would look more natural to define the operator $\partial_t^{(N)}$ – or at least the product operator $\star_K \partial_t^{(N)}$ appearing in Eq. (5.8) – by a formula which involves the (anti-symmetrized) operator $\bar{\partial}_t^{(N)}$ rather than the operators $\hat{\partial}_t^{(N)}$ and $\check{\partial}_t^{(N)}$.

And in fact, using the identification (D.4) we can easily obtain an alternative definition of the product $\star_K \partial_t^{(N)}$ which involves operators of the type "$\bar{\partial}_t$". In order to do so consider the space $\mathcal{A}^{\perp}_{2N}(K)$ and the operator $\star_K : \mathcal{A}^{\perp}_{2N}(K) \to \mathcal{A}^{\perp}_{2N}(K)$ (defined totally analogously as the space $\mathcal{A}^{\perp}(K)$ and the operator $\star_K : \mathcal{A}^{\perp}(K) \to \mathcal{A}^{\perp}(K)$ in Sec. 5.2 above but with $2N$ playing the role of N). Let $\partial_t^{(2N)} : \mathcal{A}^{\perp}_{2N}(K) \to \mathcal{A}^{\perp}_{2N}(K)$ be given by

$$\partial_t^{(2N)} := \begin{pmatrix} \bar{\partial}_t^{(2N)} & 0 \\ 0 & \bar{\partial}_t^{(2N)} \end{pmatrix} \tag{D.6}$$

where the operator $\bar{\partial}_t^{(2N)} \in \mathrm{Map}(\mathbb{Z}_{2N}, C^1(K_j, \mathfrak{g}))$, $j = 1, 2$ appearing on the main diagonal is defined totally analogously as in Definition 5.2 above (with $\mathbb{Z}_N$ replaced by $\mathbb{Z}_{2N}$).

Clearly, we can consider $\mathcal{A}^{\perp}_{altern}(K)$ as a subspace of $\mathcal{A}^{\perp}_{2N}(K)$ in the obvious way. Even though neither of the two operators $\star_K$ nor $\partial_t^{(2N)}$ leaves the subspace $\mathcal{A}^{\perp}_{altern}(K)$ of $\mathcal{A}^{\perp}_{2N}(K)$ invariant the composition $\star_K \partial_t^{(2N)} : \mathcal{A}^{\perp}_{2N}(K) \to \mathcal{A}^{\perp}_{2N}(K)$ does. It turns out that under the identification (D.4) above the restriction of $\star_K \partial_t^{(2N)}$ onto $\mathcal{A}^{\perp}_{altern}(K)$ coincides with the operator $\star_K \partial_t^{(N)} : \mathcal{A}^{\perp}(K) \to \mathcal{A}^{\perp}(K)$ appearing in Sec. 5.2 above.

Remark D.1 i) The "true meaning" (within the context of BF_3-theory) of the whole space $\mathcal{A}^{\perp}_{2N}(K)$ will become clear in Appendix E in [28].

ii) Observe that if we rewrite Sec. 5 using the space $\mathcal{A}^{\perp}_{altern}(K)$ instead of $\mathcal{A}^{\perp}(K)$ it would be convenient to undo the identification $\mathbb{Z}_N \cong \mathbb{Z}'_N$ in Sec. 4.4. Instead we identify $\mathbb{Z}_N$ with $\mathbb{Z}_{2N}^{even}$ and $\mathbb{Z}'_N$ with $\mathbb{Z}_{2N}^{odd}$. (Observe that in this case $\mathbb{Z}_{2N}$ appears naturally as the joint barycentric subdivision of $\mathbb{Z}_N$ and $\mathbb{Z}'_N$).

Moreover, we should then use Adams' original definition of "framing" in Sec. 4.4. In particular, a framing of a loop l in $\mathcal{K} \times \mathbb{Z}_N \cong K_1 \times \mathbb{Z}_{2N}^{even}$ will then be a loop l' in $\mathcal{K}' \times \mathbb{Z}'_N \cong K_2 \times \mathbb{Z}_{2N}^{odd}$.

E Continuum limits

E.1 The role of continuum limits in a purely simplicial approach

In the present series of papers we are studying the question whether it is possible to find a rigorous "discretization" approach to non-Abelian CS theory (or BF_3-theory). Ideally, such an approach would be "purely simplicial", i.e. it would not involve any kind of continuum limit. However, this does not mean that continuum limit considerations will be irrelevant if a purely simplicial approach can be found:

i) Continuum limit considerations can be useful for motivating or checking some of the constructions of a purely simplicial approach, for example by looking at the continuum limit of the RHS of Eq. (5.8) above we get an additional[65] argument for the factor 1/2 on the RHS of Eq. (5.1).

ii) For some of the possible applications of such a (hypothetical) simplicial theory, like, e.g., a new approach for proving the so-called "asymptotic expansion conjecture", cf. [4], the study of continuum limits would obviously be crucial.

E.2 Discretization approaches involving continuum limits

If a purely simplicial approach to non-Abelian CS theory (or BF_3-theory) cannot be found then discretization approaches involving a continuum limit will be the "next best thing". Of course, if we leave the simplicial setting we obviously pay a price – the use of continuum limits makes things more complicated. On the other hand we gain a certain amount of flexibility.

For example, if we work with the original discretization of the continuum operator $\frac{\partial}{\partial t} + \mathrm{ad}(B)$ appearing in Eq. (5.7) we cannot expect a relation like Eq. 5.38 (even after carrying out modifications (Mod1)–(Mod3)). However, after performing a suitable continuum limit in the S^1-component we can derive another version of Theorem 5.8, cf. Remark 5.9 above.

Similarly, by including a continuum limit "in the Σ-component" we would no longer be forced to define $A_{\mathrm{sg}}(\mathrm{h})(l^{(k)}_{\Sigma})$ by Eq. (5.15) above but we could use alternative definitions like, e.g., $A_{\mathrm{sg}}(\mathrm{h})(l^{(k)}_{\Sigma}) := A_{\mathrm{sg}}(\mathrm{h})((l^{(k)}_{\Sigma})'(0))$ or $A_{\mathrm{sg}}(\mathrm{h})(l^{(k)}_{\Sigma}) := A_{\mathrm{sg}}(\mathrm{h})((l^{(k)}_{\Sigma})'(1))$. After performing a continuum limit in the Σ-component all these different choices will lead to the same result.

Finally, let us mention that the power of continuum limits should not be overestimated. For example, at first look one could think that when working with a continuum limit in the Σ-component, it does not matter anymore whether we use the original definition of $B(\bar{e})$ as in Eq. (5.10) above or the modified definition Eq. (5.35) given in (Mod2). However, a closer look shows that the explicit values of the various "continuum limit versions" of $\mathrm{WLO}_{rig}(L)$ indeed depend on the precise definition of $B(\bar{e})$ which we use.

---

[65] in case part D of the Appendix has not yet convinced the reader of this point

F Appendix: A comment on the "continuum approach" towards a rigorous realization of the RHS of Eq. (2.53)

In Secs. 1–2 we sometimes referred to the "continuum approach" of [24, 26, 27] for making rigorous sense of the heuristic expression for WLO(L) appearing[66] on the RHS of Eq. (2.53) This approach is based on white noise analysis (WNA) (cf. [31]) and was in part inspired by [3, 23]. For the convenience of the reader we will now give a brief summary of the main aspects of this approach.

i) Use of WNA: Consider the heuristic integral functionals associated to the two heuristic Gauss-type complex measures appearing in Eq. (2.53) in Sec. 2.3.3, i.e.

$$\Phi^{\perp}_{B} := \int \cdots \exp(iS_{CS}(\hat{A}^{\perp}, B)) D\hat{A}^{\perp} \tag{F.1a}$$

$$\Psi := \int \cdots \exp(iS_{CS}(A^{\perp}_c, B))(DA^{\perp}_c \otimes DB) \tag{F.1b}$$

We would like to find a rigorous realization of $\Phi^{\perp}_{B}$ as a continuous linear functional on a certain infinite dimensional function space. (Similar considerations can be made for the functional Ψ.)

Note that even if $\exp(iS_{CS}(\hat{A}^{\perp}, B))D\hat{A}^{\perp}$ were a (heuristic) probability measure one could not hope to realize $\Phi^{\perp}_{B}$ rigorously as a functional $C_b(\hat{\mathcal{A}}^{\perp}, \mathbb{R}) \to \mathbb{C}$, cf. Remark F.1 i) below. In order to succeed one would have to replace $C_b(\hat{\mathcal{A}}^{\perp}, \mathbb{R})$ by $C_b(\overline{\hat{\mathcal{A}}^{\perp}}, \mathbb{R})$ where $\overline{\hat{\mathcal{A}}^{\perp}}$ is a suitable extension of $\hat{\mathcal{A}}^{\perp}$.

For an oscillatory Gauss-type measure like $\exp(iS_{CS}(\hat{A}^{\perp}, B))D\hat{A}^{\perp}$ one can not even hope to be able to define rigorously as a continuous linear functional on the full space $C_b(\overline{\hat{\mathcal{A}}^{\perp}}, \mathbb{R})$ but only on a suitable subspace $\mathcal{X}$ of $C_b(\overline{\hat{\mathcal{A}}^{\perp}}, \mathbb{R})$ or $C(\overline{\hat{\mathcal{A}}^{\perp}}, \mathbb{R})$, cf. Remark F.1 ii) below. The framework of WNA provides a general method for constructing such spaces. Using this method for the construction of a suitable space[67] $\mathcal{X}$ we obtained in [24, 26] a rigorous realization of $\Phi^{\perp}_{B}$ as a continuous linear functional $\Phi^{\perp}_{B} : \mathcal{X} \to \mathbb{C}$.

ii) "Loop smearing": In point i) we sketched how one can realize $\Phi^{\perp}_{B}$ rigorously as a continuous linear functional $\mathcal{X} \to \mathbb{C}$ on a certain topological space $\mathcal{X}$ appearing as a linear subspace of $C_b(\overline{\hat{\mathcal{A}}^{\perp}}, \mathbb{R})$. Since the extension $\overline{\hat{\mathcal{A}}^{\perp}}$ of $\hat{\mathcal{A}}^{\perp}$ contains non-smooth elements $\hat{A}^{\perp}$, expressions like $\hat{A}^{\perp}(l'(t))$ and $\mathrm{Tr}_{\rho_i}\big(\mathrm{Hol}_{l_i}(\hat{A}^{\perp} + A^{\perp}_c, B; \mathrm{h})\big)$ are not defined for general $\hat{A}^{\perp} \in \overline{\hat{\mathcal{A}}^{\perp}}$, cf. Eq. (2.4). This complication can be resolved by using "loop smearing", which is analogous to "point smearing" described in Remark F.1 i) below.

iii) "Framing": In all (heuristic or rigorous) approaches for making sense of the (Abelian or Non-Abelian) CS path integral some form of "framing" has to be used. For the implementation of the framing procedure in [24, 26, 27] we fixed a suitable family $(\phi_s)_{s>0}$ of diffeomorphisms of $M = \Sigma \times S^1$ fulfilling $(\phi_s)^* \mathcal{A}^{\perp} = \mathcal{A}^{\perp}$ for all $s > 0$ and $\phi_s \to \mathrm{id}_M$ as $s \to 0$ uniformly (w.r.t. to an arbitrary Riemannian metric on $M = \Sigma \times S^1$).

For each diffeomorphism $\phi \in \{\phi_s \mid s > 0\}$ one can introduce a suitable "deformation" $S_{CS,\phi}(A^{\perp} + Bdt)$ of $S_{CS}(A^{\perp} + Bdt)$ (cf. Eq. (F.2) below) and a corresponding deformation $\Phi^{\perp}_{B,\phi} := \int \cdots \exp(iS_{CS,\phi}(\hat{A}^{\perp}, B))D\hat{A}^{\perp}$ of the integral functional $\Phi^{\perp}_{B}$ above. (And in a similar way one can introduce a deformation Ψ_{ϕ} of Ψ).

---

[66] or, more precisely, the version of Eq. (2.53) which one obtains when working with the decomposition $\mathcal{A}^{\perp} = \hat{\mathcal{A}}^{\perp} \oplus \mathcal{A}^{\perp}_c$ instead of $\mathcal{A}^{\perp} = \check{\mathcal{A}}^{\perp} \oplus \mathcal{A}^{\perp}_c$

[67] The space $\mathcal{X}$ was denoted by $(\mathcal{N})$ in [24, 26] where $\mathcal{N} := \mathcal{A}^{\perp}$ equipped with a suitable family of seminorms.

In [24, 26, 27] we implicitly[68] worked with $S_{CS,\phi}(A^{\perp} + Bdt)$ given by

$$S_{CS,\phi}(A^{\perp} + Bdt) = \pi k \big[\ll \phi^* A^{\perp}, \star\big(\tfrac{\partial}{\partial t} + \operatorname{ad}(B)\big) A^{\perp} \gg -2 \ll \phi^* A^{\perp}, \star dB \gg \big] \tag{F.2}$$

iv) ζ-function regularization: In [11] the heuristic expression

$$Det(B) := \det\big(1_{\mathfrak{k}} - \exp(\operatorname{ad}(B))_{|\mathfrak{k}}\big)\hat{Z}(B) \tag{F.3}$$

where $\hat{Z}(B) := \int \exp(iS_{CS}(\hat{A}^{\perp}, B))D\hat{A}^{\perp}$ was evaluated explicitly for constant B in two different ways, cf. Sec. 3 and Sec. 6 in [11]. Both evaluations involve a suitable ζ-function regularization[69]. At a formal level the formula in [11] can immediately be generalized to the case where B is a step function, cf. Sec. 3.5 in [26]. Let us remark, however, that a careful justification of the general formula for $Det(B)$ has not been given yet (cf. Remark 3.3 in [26] and Remark 4.2 in [25]).

The following remark is relevant for point i) in the list above.

Remark F.1 i) Consider the pre-Hilbert space $\mathcal{F} := (C^{\infty}_{\mathbb{R}}(\mathbb{R}), \langle \cdot, \cdot \rangle)$ where $\langle \cdot, \cdot \rangle$ is the scalar product induced by $L^2_{\mathbb{R}}(\mathbb{R}, dx)$ and let $S : \mathcal{F} \to \mathcal{F}$ be a positive invertible linear operator. Let us consider the heuristic integral

$$\int_{\mathcal{F}} \prod\nolimits_{i=1}^{n} f(x_i) d\mu_S(f) \tag{F.4}$$

where $(x_i)_{i \leq n}$ is a fixed sequence of points in $\mathbb{R}$ and where $d\mu_S$ is the heuristic Gaussian measure on $\mathcal{F}$ given by

$$d\mu_S(f) = \exp(-\langle f, Sf \rangle)df \tag{F.5}$$

df being the heuristic Lebesgue measure on $\mathcal{F}$.

It turns out that in general it is not possible to make rigorous sense of the heuristic measure $d\mu_S$ as a Borel measure on $\mathcal{F}$. However, one can make sense of this heuristic measure as a Borel measure on a suitable extension $\overline{\mathcal{F}}$ of $\mathcal{F}$, for example on $\overline{\mathcal{F}} := \mathcal{S}'_{\mathbb{R}}(\mathbb{R})$ (equipped with the standard topology).

Even though we then have a rigorous realization of $d\mu_S$ it is still not possible to make sense of the heuristic expression (F.4) without further steps. This is because for a general element f of $\mathcal{S}'_{\mathbb{R}}(\mathbb{R})$ the expression $f(x_i)$ does not make sense so the integrand in Eq. (F.4) is ill-defined. This complication can often be resolved by "point smearing", i.e. in Eq. (F.4) we replace, for each $\epsilon > 0$, the "point" x_i by a test function $\phi^{\epsilon}_i \in \mathcal{S}(\mathbb{R})$ such that $\phi^{\epsilon}_i \geq 0$, $\int \phi^{\epsilon}_i(x)dx = 1$ and $\operatorname{supp}(\phi^{\epsilon}_i)$ is contained in a small ϵ-neighborhood of x_i. The parameter ϵ is later sent to 0.

ii) For every bounded complex Borel measure $d\mu$ on a topological space Y the associated integral functional $\int \cdots d\mu$ is a continuous linear functional $C_b(Y, \mathbb{R}) \to \mathbb{C}$. Clearly, for unbounded complex Borel measures the corresponding integral functional $\int \cdots d\mu$ can in general not be defined as a linear functional on the whole space $C_b(Y, \mathbb{R})$ but only on a suitable subspace $\mathcal{X}$ of $C_b(Y, \mathbb{R})$ or, more generally, a "sufficiently small" subspace of $C(Y, \mathbb{R})$. For example, in the special case where $Y = \mathbb{R}^n$ with the standard scalar product and where $d\mu$ is an oscillatory Gauss-type measure on $\mathbb{R}^n$ convenient choices for $\mathcal{X}$ are $\mathcal{X} = \mathcal{S}(\mathbb{R}^n)$ or $\mathcal{X} = \mathcal{P}_{exp}(\mathbb{R}^n)$, cf. Sec. 3.

---

[68] in fact, the explicit heuristic formulas which actually appeared in [24, 26, 27] are those describing the informal Fourier-transform of the integral functional $\Phi^{\perp}_{B,\phi}$, This is because for the rigorous realization of $\Phi^{\perp}_{B,\phi}$ we applied the Kondratiev-Potthoff-Streit characterization theorem of White noise analysis (an analogue of the Bochner-Minlos theorem)

[69] additionally, the first evaluation involves an argument based on Ray-Singer torsion and the second evaluation uses an index theorem

References

[1] D. H. Adams. R-torsion and linking numbers from simplicial abelian gauge theories. Preprint, arXiv:hep-th/9612009, 1996.

[2] D. H. Adams. A doubled discretization of abelian Chern-Simons theory. *Phys. Rev. Lett.*, 78(22):4155–4158, 1997.

[3] S. Albeverio and A.N. Sengupta. A Mathematical Construction of the Non-Abelian Chern-Simons Functional Integral. *Commun. Math. Phys.*, 186:563–579, 1997.

[4] J.E. Andersen. The asymptotic expansion conjecture, Sec. 7.2. in *Problems on invariants of knots and 3-manifolds*, ed. T. Ohtsuki, in *Invariants of knots and 3-manifolds*, Geometry and Topology Monographs 4 (2002), 377–572.

[5] M. Atiyah. *The Geometry and Physics of Knots*, Cambridge U. Press, Cambridge, 1990.

[6] John C. Baez. Four-dimensional BF theory as a topological quantum field theory. *Lett. Math. Phys.*, 38(2):129–143, 1996.

[7] J. Barrett and I. Naish-Guzman. The Ponzano-Regge model, arXiv:0803.3319

[8] C. Beasley and E. Witten. Non-abelian localization for Chern-Simons theory. *J. Differential Geom.*, 70(2):183–323, 2005.

[9] C. Beasley. Localization for Wilson Loops in Chern-Simons Theory. arXiv:0911.2687, 2009

[10] M. Blau and G. Thompson. A new class of topological field theories and the Ray-Singer torsion. *Phys. Lett. B*, 228(1):64–68, 1989.

[11] M. Blau and G. Thompson. Derivation of the Verlinde Formula from Chern-Simons Theory and the G/G model. *Nucl. Phys.*, B408(1):345–390, 1993.

[12] M. Blau and G. Thompson. Lectures on 2d Gauge Theories: Topological Aspects and Path Integral Techniques. In E. Gava et al., editor, *Proceedings of the 1993 Trieste Summer School on High Energy Physics and Cosmology*, pages 175–244. World Scientific, Singapore, 1994.

[13] M. Blau and G. Thompson. On Diagonalization in $Map(M, G)$. *Commun. Math. Phys.*, 171:639–660, 1995.

[14] M. Blau and G. Thompson. Chern-Simons theory on S^1-bundles: abelianisation and q-deformed Yang-Mills theory. *J. High Energy Phys.*, (5):003, 35 pp. (electronic), 2006.

[15] Th. Bröcker and T. tom Dieck. *Representations of compact Lie groups*, volume 98 of *Graduate Texts in Mathematics.* Springer-Verlag, New York, 1985.

[16] A. Cattaneo, P. Cotta-Ramusino, J. Fröhlich, and M. Martellini. Topological BF theories in 3 and 4 dimensions. *J. Math. Phys.*, 36(11):6137–6160, 1995.

[17] J. Cheeger. Analytic torsion and the heat equation. *Ann. of Math. (2)*, 109(2):259–322, 1979.

[18] S. de Haro and A. Hahn. Chern-Simons theory and the quantum Racah formula. Preprint, arXiv:math-ph/0611084.

[19] D. S. Fine. On Chern-Simons and WZW partition functions. *Comm. Math. Phys.*, 200(3):685–698, 1999.

[20] L. Freidel and D. Louapre. Ponzano-Regge model revisited. I. Gauge fixing, observables and interacting spinning particles. *Classical Quantum Gravity*, 21(24):5685–5726, 2004.

[21] L. Freidel and D. Louapre. Ponzano-Regge model revisited II: Equivalence with Chern-Simons. Preprint, arXiv:gr-qc/0410141v3, 2004.

[22] L. Freidel and E. R. Livine. Ponzano-Regge model revisited. III. Feynman diagrams and effective field theory. *Classical Quantum Gravity*, 23(6):2021–2061, 2006.

[23] A. Hahn. The Wilson loop observables of Chern-Simons theory on $\mathbb{R}^3$ in axial gauge. *Commun. Math. Phys.*, 248(3):467–499, 2004.

[24] A. Hahn. Chern-Simons models on $S^2 \times S^1$, torus gauge fixing, and link invariants I. *J. Geom. Phys.*, 53(3):275–314, 2005.

[25] A. Hahn. Chern-Simons models on $S^2 \times S^1$, torus gauge fixing, and link invariants II. *J. Geom. Phys.*, 58:1124–1136, 2008.

[26] A. Hahn. An analytic Approach to Turaev's Shadow Invariant. *J. Knot Th. Ram.*, 17(11): 1327–1385, 2008 [see arXiv:math-ph/0507040v7 (2011) for the most recent version]

[27] A. Hahn. White noise analysis in the theory of three-manifold quantum invariants. In A.N. Sengupta and P. Sundar, editors, *Infinite Dimensional Stochastic Analysis*, volume XXII of *Quantum Probability and White Noise Analysis*, pages 201–225. World Scientific, 2008.

[28] A. Hahn. From simplicial Chern-Simons theory to the shadow invariant II, Preprint, 2012.

[29] A. Hahn, B. McLellan, and B. Himpel, in preparation.

[30] A. Hatcher. *Algebraic Topology.* Cambridge U. Press, Cambridge, 2002.

[31] T. Hida, H.-H. Kuo, J. Potthoff, and L. Streit. *White Noise. An infinite dimensional Calculus.* Dordrecht: Kluwer, 1993.

[32] S.-T. Hu. *Homotopy Theory.* Academic Press, New York and London, 1959.

[33] Y. Katznelson. *An introduction to harmonic analysis*, *Harmonic Analysis*, Cambridge University Press, 2004

[34] A. N. Kirillov and N.Y. Reshetikhin. Representations of the algebra $U_q(sl_2)$, q-orthogonal polynomials and invariants of links. In V.G. Kac et al., editor, *Infinite Dimensional Lie Algebras and Groups*, Vol. 7 of *Advanced Ser. in Math. Phys.*, pages 285–339, 1988.

[35] D. Krotov and A. Losev. Quantum Field Theory as Effective BV Theory from Chern-Simons. arXiv:hep-th/0603201.

[36] G. Kuperberg. Quantum invariants of knots and 3-manifolds (book review). *Bull. Amer. Math. Soc.*, 33(1):107–110, 1996.

[37] J. M. F. Labastida. Chern-Simons Gauge Theory: ten years after, arXiv: hep-th/9905057v1

[38] P. Mnëv. Notes on simplicial BF theory, arXiv: hep-th/0610236v3

[39] W. Müller. Analytic torsion and R-torsion of Riemannian manifolds. *Adv. in Math.*, 28(3):233–305, 1978.

[40] S. Pokorski. *Gauge Field Theories.* Cambridge U. Press, Cambridge, 2000.

[41] N.Y. Reshetikhin and V.G. Turaev. Ribbon graphs and their invariants derived from quantum groups. *Commun. Math. Phys.*, 127:1–26, 1990.

[42] N.Y. Reshetikhin and V.G. Turaev. Invariants of three manifolds via link polynomials and quantum groups. *Invent. Math.*, 103:547–597, 1991.

[43] J. Roberts. Skein theory and Turaev-Viro invariants. *Topology*, 34(4):771–787, 1995.

[44] L. Rozansky. A contribution of the trivial connection to Jones polynomial and Witten's invariant of 3d manifolds. I, arXiv:hep-th/9401061

[45] S. F. Sawin. Jones-Witten invariants for nonsimply connected Lie groups and the geometry of the Weyl alcove, *Advances in Mathematics*, 165(1):1–34, 2002 [arXiv:math/9905010v2]

[46] S. F. Sawin. Quantum groups at roots of unity and modularity. *J. Knot Theory Ramifications*, 15(10):1245–1277, 2006.

[47] V. G. Turaev. *Quantum invariants of knots and 3-manifolds.* de Gruyter, 1994.

[48] V.G. Turaev. Shadow links and face models of statistical mechanics. *J. Diff. Geom.*, 36:35–74, 1992.

[49] E. Witten. Quantum Field Theory and the Jones Polynomial. *Commun. Math. Phys.*, 121:351–399, 1989.